

World Journal of *Clinical Infectious Diseases*

World J Clin Infect Dis 2023 May 29; 13(2): 11-23



MINIREVIEWS

- 11 Emerging leishmaniasis in southern Himalayas: A mini-review

Sharma A, Kumar S, Panda PK, Yadav S, Kalita D

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Infectious Diseases*, Ahmed Morad Asaad, MD, Professor, Microbiology Department, College of Medicine, Zagazig University, Zagazig 44519, Egypt.
ahmedmoradasaad@hotmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Infectious Diseases* (WJCID, *World J Clin Infect Dis*) is to provide scholars and readers from various fields of infectious diseases with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJCID mainly publishes articles reporting research results and findings obtained in the field of infectious diseases and covering a wide range of topics including community-acquired infections, cross infection, eye infections, focal infection, infectious gingivitis, intraabdominal infections, laboratory infection, Ludwig's angina, necrotizing ulcerative periodontitis, opportunistic infections, pelvic infection, pregnancy complications, etc.

INDEXING/ABSTRACTING

The WJCID is now abstracted and indexed in Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Jia-Le Jin; Production Department Director: Xu Guo; Editorial Office Director: Yun-Xiao Jiao Wu.

NAME OF JOURNAL

World Journal of Clinical Infectious Diseases

ISSN

ISSN 2220-3176 (online)

LAUNCH DATE

December 30, 2011

FREQUENCY

Continuous Publication

EDITORS-IN-CHIEF

Joao Mesquita, Caterina Sagnelli, Wei Wang, Haroon Ahmed

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2220-3176/editorialboard.htm>

PUBLICATION DATE

May 29, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/gerinfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/gerinfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/gerinfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Emerging leishmaniasis in southern Himalayas: A mini-review

Ashwani Sharma, Santosh Kumar, Prasan Kumar Panda, Sweetly Yadav, Deepjyoti Kalita

Specialty type: Infectious diseases

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C, C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Mukhopadhyay S, United States; Rychtář J, United States

Received: September 28, 2022

Peer-review started: September 28, 2022

First decision: November 22, 2022

Revised: November 28, 2022

Accepted: April 27, 2023

Article in press: April 27, 2023

Published online: May 29, 2023



Ashwani Sharma, Santosh Kumar, Department of Community and Family Medicine, All India Institute of Medical Sciences, Rishikesh 249203, India

Prasan Kumar Panda, Department of Medicine, All India Institute of Medical Sciences, Rishikesh 249203, India

Sweetly Yadav, Department of Internal Medicine, All India Institute of Medical Sciences, Rishikesh 249203, India

Deepjyoti Kalita, Department of Microbiology, All India Institute of Medical Sciences, Rishikesh 249203, India

Corresponding author: Prasan Kumar Panda, MBBS, MD, Associate Professor, Department of Medicine, All India Institute of Medical Sciences, Sixth Floor, College Block, Rishikesh 249203, India. motherprasanna@rediffmail.com

Abstract

Leishmaniasis is a vector-borne parasitic disease affecting millions of people worldwide. However, in the last decade, the number of cases has been reduced from well-documented endemic parts, but sporadic cases have been reported widely from various non-endemic areas, especially from the southern Himalayan zone. This raises concerns about the emergence of new ecological niches. This warrants a critical evaluation of key factors causing this rapid spread and possibly indigenous transmission. This mini-review article is aimed to briefly address the parasite, the vector, and the environmental aspects in the transmission of leishmaniasis in these new foci against a background of worldwide endemic leishmaniasis with a special focus on the southern Himalayan zone. As the lack of knowledge about the causative parasites, vectors, reservoir hosts, atypical presentations, and their management make the problem serious and may lead to the emergence of public health issues. The present work also reviewed the existing information regarding clinical variations, diagnostic methods, treatment, its outcome, and ignite for further research in these aspects of the disease.

Key Words: Anthroponosis; Kala azar sandfly; Sporadic transmission; Southern Himalaya

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This mini-review article is aimed to briefly address the parasite, the vector, and the environmental aspects in the transmission of leishmaniasis in these new foci against a background of worldwide endemic leishmaniasis with a special focus on the southern Himalayan zone.

Citation: Sharma A, Kumar S, Panda PK, Yadav S, Kalita D. Emerging leishmaniasis in southern Himalayas: A mini-review. *World J Clin Infect Dis* 2023; 13(2): 11-23

URL: <https://www.wjgnet.com/2220-3176/full/v13/i2/11.htm>

DOI: <https://dx.doi.org/10.5495/wjcid.v13.i2.11>

INTRODUCTION

Leishmaniasis is a vector-borne parasitic disease that exists either as zoonosis (in most endemic parts of the world) or anthroponosis (endemic part of the Indian sub-continent) and is transmitted by S and fly. The latter entity is on the verge of elimination, efficiently with the help of assigned memorandum of understanding by the five most endemic countries: India, Nepal, Bangladesh, Bhutan, and Thailand[1]. But at present, a major challenge is the increasing emergence of new ecological niches having indigenous transmission. Recently World Health Organization (WHO) declares it as a category I disease (emerging and uncontrolled), and the World Health Assembly recognizes it as a major public health concern[2]. Leishmaniasis is a disease of low altitude. It does not occur at an altitude of more than 2000ft (600m)[3]. The Southern Himalayan regions (of countries like Pakistan, India, Nepal, and Bhutan) are considered as non-endemic regions probably because of the non-conducive environment for the growth of its vector, *i.e.*, Sand flies. But as several cases of leishmaniasis have been reported from the sea areas, the above observational facts are being indistinct. Most of these cases were found along with the upstream of Himalayas river belts (like Indus, Ganga, Yamuna, and the Brahmaputra) especially in the western part (Islamabad, Jammu & Kashmir, and Himachal Pradesh), the middle part (Uttarakhand), and the eastern part (Nepal and Bhutan) of Himalayas[3-33].

Here, a mini-/narrative review is done considering the available case reports/case series/observational studies from new emerging areas, regarding leishmaniasis disease profile (epidemiology, microbiology, patho-physiology, clinical variations, diagnostic methods, treatment, and outcome, including the entomological assessment of S and fly). This article also intends to focus on the difference between the disease profile of leishmaniasis in the southern Himalayan belt *vs* the world's endemic areas in a systematic manner.

METHODOLOGY

A mini-review of all published (PubMed/Medline, Embase, Cochrane database, Google Scholar) leishmania cases from the Himalayas regions of Pakistan, India, Nepal, and Bhutan were reviewed and analyzed with prime focus on the disease profiles of the cases reported in the lower Himalayan belt (Figure 1). For distinctive comparison and obtaining good inference, the Indian Himalayan belt is further divided into Jammu & Kashmir, Himachal Pradesh (Shimla, Chamba & Kinnaur), and Uttarakhand (Garhwal & Kumaon) regions. Leishmaniasis which was initially considered a disease of plain lower altitude areas along the banks of major rivers is now prevailing in higher altitudes. This ecological shift provides us with an excellent opportunity to study the epidemiological triad and also warranting a need to implement appropriate control measures. Hence, this review is done with the objective to identify the newly reported endemic areas on these hilly terrains related to the disease and multiple factors associated with it, especially in relation to river belts.

Selection: (1) Leishmania disease: Only records that concern the leishmania/Kala-azar in the Indian sub-continent or related topics are included in the selection; (2) Original records: We excluded letters, editorials and comments; and (3) English language: We excluded articles written in other language.

RESULTS

Across all literature and records available, 31 references were found which were relevant to our study (Table 1) among 51 qualitative syntheses (Figure 2). The sample size in these studies varied from a single case report to a study containing more than thousands of cases[3-33].

Epidemiology (demography)

The studies reviewed were specially chosen from the southern Himalayan region to emphasize the

Table 1 Findings of published articles on leishmaniasis in southern Himalayas w.r.t. agent-host-environment details

Ref.	Sample size	Location (if available district, state, country)	Agent factors	Host factors	Vector identified	Environmental factors	River body associated	Authors conclusion
Katakura <i>et al</i> [4]	> 1000	Different areas of Pakistan, India, and Nepal	In Pakistan Himalayas, <i>Leishmania tropica</i> followed by <i>Leishmania major</i>	CP: Cutaneous leishmaniasis (CL) cases only; No descriptions Rx: Not known	In all Himalayas, <i>P. sergenti</i> followed by <i>P. argentipes</i> and <i>papatasi</i>	Altitude is not documented	Indus, Ganges	Microsatellite analysis of the parasites will be a powerful tool for population genetic and epidemiological studies of <i>Leishmania</i> species
			In India, <i>L. donovani</i> followed by <i>L. tropica</i>	Outcome: Not known				
			In Nepal, <i>L. major</i>					
Rab <i>et al</i> [5]	239 (1984-1992) and more cases in the past (before 1984)	Different areas of Northern areas of Pakistan (Bagh, Abbottabad, Chilas, and Baltistan)	<i>Leishmania</i>	Clinical presentation (CP): <i>Visceral leishmaniasis</i> (VL – all cases); Not described	Not documented	Altitude is not documented	Indus	The clinical pattern of VL in north Pakistan is akin to that in north-western China, with a marked predilection for young children, and a male preponderance. The infantile VL has risen 10-fold in the last decade from 0.2 to almost 2 per 100 000 population
			<i>infantum</i>	Rx: Not known Outcome: Not known				
Wani <i>et al</i> [6]	18	Different areas of Uri & Karnah belt, Jammu & Kashmir, India	<i>Leishmania</i> , species not identified	CP: Cutaneous leishmaniasis (CL); mostly nodulo-ulcerative, mostly on the face and single lesion Rx: Intravenous sodium Stibogluconate including two received intra-lesional Outcome: Survival for all cases	Not documented	Altitude is not documented. The hot and arid climate of these areas (Uri belt) is quite conducive to the growth and development of leishmania and the sand fly	Not documented	Any patient with nodular/nodulo-ulcerative lesion on exposed parts must be suspected for CL, especially if belonging to the Uri and Karnah region of the Kashmir Valley. The public health authorities should make every effort to contain this new infection in this Valley
Leherwal <i>et al</i> [7]	Three	Uri belt, Jammu & Kashmir, India	<i>Leishmania</i> , species not identified	CP: Cutaneous leishmaniasis (CL); solitary erythematous nodule on the face Rx: Not documented Outcome: Not documented	Not documented	Altitude is not documented	-do-	Focuses on the diagnostic part. FNAC may be the method of choice for suspected CL in cases of solitary nodular lesions
Mahajan <i>et al</i> [8]	One	Uri in South West Kashmir, Jammu & Kashmir, India	<i>Leishmania</i> , species not identified	CP: Visceral leishmaniasis (VL); 2 months fever, weight loss, ascites, anemia, Hepato-splenomegaly, Rx: Intravenous sodium	Not documented	Altitude is not documented	-do-	This advice for further research into the epidemiology, geographic distribution, and inter-species interactions of the parasite

Sharma <i>et al</i> [9]	285	Nirmand village, Shimla & Kullu Districts of Himachal Pradesh, India	Among 14 cases, <i>Leishmania tropica</i> (3) and <i>Leishmania donovani</i> (11)	Stibogluconate	Among 41 cases, <i>P. longiductus</i> (29), <i>P. major</i> (8), <i>P. kandelaki</i> (2), and 2 remained unidentified	Altitude is not documented	Satluj river	Different leishmania species and vectors compared to other parts of India are found in these Himalayas
				Outcome: Survived				
Sharma <i>et al</i> [10]	161 new localized cases of LCL from May 2001 and December 2003	sub-alpine valley in the mountainous region of the Kinnaur District, Himachal Pradesh, India	<i>L. donovani</i> in eight cases and <i>L. tropica</i> in two cases	CP: CL; mostly nodulo-ulcerative, mostly on extremities	<i>Phlebotomus longiductus</i> is a possible vector	Altitude, 700-2,900 m above sea level	Satluj River	Intralesional sodium stibogluconate was effective in all patients
				Tissue smear positivity for amastigotes was 43%				
Raina <i>et al</i> [11]	18	Shimla, Kinnaur & Kullu Districts of Himachal Pradesh, India	Leishmania, species not identified	Rx: Intra-leisonal sodium	Not documented	Altitude, 924 - 2960 m above sea level	Satluj and Beas river	Initial failure to suspect VL in this area might cause a diagnostic delay
				Stibogluconate				
Thakur <i>et al</i> [46]	Cases of CL During 2014–2018 in the study area	case reports came from Districts of Kinnaur, Shimla, and Kullu and the previously nonendemic districts of Mandi and Solan, Himachal Pradesh, India	<i>L. donovani</i> variants distinct from the viscerotropic <i>L. donovani</i> strain from northeast India	Outcome: Survival for all cases	Coexistence of VL and CL	Not documented	Not documented	The scenario appears somewhat similar to Sri Lanka and Kerala, where <i>L. donovani</i> parasites cause cutaneous disease, albeit with differences in the region-specific <i>L. donovani</i> variants
				Histopathology showed non-caseating epitheloid cell granuloma in 77% of the cases. Lesions involved mainly the face				
Thakur <i>et al</i> [47]	Sixty CL patients over the period from 2014 to 2018	Satluj river belt in Himachal Pradesh, Khaneri/rampur (location of medical college), Himachal Pradesh, India	Presence of <i>L. seymouri</i> co-infection in the unusual CL cases in Himachal Pradesh (HP) caused by <i>L. donovani</i> variants	Rx: Intravenous sodium	Coexistence of VL and CL	Not documented	Not documented	Found the presence of <i>Leptomonas seymouri</i> in 38.5% (22/57) of the patients along with <i>L. donovani</i> detected in all the samples. <i>L. seymouri</i> is a monoxenous insect trypanosoma, generally incapable of infecting humans
				Stibogluconate				
				Outcome: 14 Survives and 4 deaths		The patients had never visited any of the endemic areas		There is a favorable therapeutic response without recurrence of symptoms during 6 months of follow-up

Sharma <i>et al</i> [49]	None	Shimla, Kinnaur, & Kullu Districts of Himachal Pradesh, India	Not applicable	Not applicable	Among 62 cases, <i>Phlebotomus longiductus</i> (46), <i>P. major</i> (8), <i>P. kandelaki</i> (8)	Our patients reported having been out of the state or district during the three years the preceding onset of symptoms	Satluj river	<i>Phlebotomus longiductus</i> may be the primary vector for human leishmaniasis in this endemic focus, however, it needs another study to prove the vector species corresponding to the type of leishmania species
--------------------------	------	---	----------------	----------------	---	---	--------------	---

growing concern of leishmania in newly endemic areas. Among all reviewed studies, one was conducted in north Pakistan, twenty-three in north India, four in Nepal, and two in Bhutan. Among Indian studies, three were in Kashmir, eleven in Himachal Pradesh (Shimla, Chamba & Kinnaur), and nine in Uttarakhand (Garhwal & Kumaon). One study was multi-centric, covering vast geographical areas falling in Pakistan, India, and Bhutan[4].

Considering the pivotal role of the environment in the natural history of disease meticulous scrutiny of various articles was done. The majority of studies included in this review have been conducted along the banks of major river-belts of the terrain (Figure 3). In northern Pakistan, the major river associated was Indus and its tributaries. In northern India, the Uri Belt of Jammu & Kashmir, the river belt of Satluj and Ravi in Himachal Pradesh, and the bank of the river Ganges in Uttarakhand were the major site of focus. In Nepal, a total of four studies have been reported which were conducted along the banks of river Budhi Ganga and Kailash. One study from the mid-west region of Nepal has not documented an associated river, but further search for location indicates the site belongs to the banks of river Karnali. Similarly, studies from eastern Bhutan have not specified associated rivers but the described areas are mainly located between the three major rivers-Drangme Chhu, Kuru Chhu, and Mangde Chhu, all are tributaries of the Brahmaputra river. A multi-national study from South and South-east Asia also reported Indus and the Ganges to be the major associated river[4]. Among all these studies none of them established a direct association between the presence of any major water bodies & ecological niche conducive for the vector species.

Entomological and parasitological findings (Table 1)


Although the majority of the reviewed studies did not identify the vector species, *Phlebotomus argentipes* was the pre dominant vector species among all the reported cases[5-7]. Few studies have also found some different species as a possible vector such as *P. longiductus* > *P. major* > *P. kandelaki* as a leading species of the vector in studies of Shimla & Kullu districts of Himachal Pradesh, India[8,9]. Similarly, one study from Bhutan has also reported four different phlebotomine species[10].

The existence of *L. donovani* was ubiquitous however the quest to identify the predominant causative leishmania species remains unresolved as the majority of the studies did not identify any. Among the studies included in our review, five studies have reported *L. donovani*[10-15], while two studies reported *L. infantum*[16], as the predominant leishmaniasis species. Few studies indicated the presence of dual-species like both *L. tropica* and *L. donovani*[4,8,9], were documented in three studies and both *L. infantum* and *L. donovani*[17] were documented in a single study. It is also recorded that *L. donovani* variants found in Himachal Pradesh, India were different from the viscera tropic leishmania strain predominant in north east India[11].



DOI: 10.5495/wjcid.v13.i2.11 Copyright ©The Author(s) 2023.

Figure 1 Map of the southern Himalayas describing rivers.



DOI: 10.5495/wjcid.v13.i2.11 Copyright ©The Author(s) 2023.

Figure 2 Map of the southern Himalayas representing the magnitude of leishmaniasis.

Clinical presentation (Table 1)

The majority of the studies reported cases of visceral leishmaniasis (VL) with high-grade prolonged fever, malaise, abdominal discomfort [3,7,10,13,14,16,18-27]. Cutaneous leishmaniasis (CL) was reported in a few studies with clinical presentation of nodulo-ulcerative lesions or solitary erythematous nodule




Figure 3 Preferred Reporting Items for Systematic reviews and Meta-Analyses flow diagram.

[4,8,9,12,28,29]. Three studies reported cases with both types (VL and CL) of leishmaniasis[11,17,30]. Another three studies did not identify the type of leishmaniasis however they described a clinical picture of hepatomegaly and weight loss as a common feature in their studies[15,31,32].

Laboratory diagnosis (Table 1)

Methods of laboratory diagnosis were not documented in any of the reviewed literature, however, smear-positive by Giemsa or Leishman technique for *Leishmania donovani* (LD) bodies are reported in most cases. LD bodies were demonstrated in the bone marrow in the case of VL and from the skin in the case of CL[13,18,28]. Some studies also found LD bodies in splenic aspirate, lymph node aspirate, duodenal and colonic mucosal biopsy in patients presenting with diarrhea[3,6,9,22]. Only in a few reference studies, there were records of other methods (mostly rK39 ICT) as an additional test. One case report of a pregnant lady was found rK16 test positive, rather than commonly used rK39 antigen[21]. Secondary hemophagocytosis lymphocytic syndrome (HLH) in VL cases was diagnosed either by 4 out of 6 criteria of HLH diagnosis or by bone marrow aspirate examination for hemophagocytosis[20,21,23]. Rarely polymerase chain reaction (PCR) for the leishmania kinetoplast mini circle gene was tested and found to be positive in a case of *L. donovani* infection which was confirmed on subsequent sequencing of the PCR – amplification method[27]. An age-old aldehyde test was found positive for five out of six cases of kala-azar, however, they confirm edit either by rK39 testing or by bone marrow aspiration examination for the L D bodies[33].

Treatment and outcome (Table 1)

Pharmacological therapies with sodium stibogluconate, amphotericin-B or miltefosine, either single or in various combinations had been reported in 21 reviewed studies. Studies were done in northern Pakistan and the Uri belt of Kashmir did not document the pharmacotherapy used and hence the subsequent outcomes[4,16,29]. In the case of VL, studies had reported intravenous sodium stibogluconate alone is sufficient for upto 84% of cases (19 survivals and 5 deaths, out of 24 cases)[8,18,25]. However, some studies were not clear about the route of stibogluconate therapy (intravenous or intralesional). Plain amphotericin-B showed > 90% recovery rate and liposomal showed upto 100% cure rate[7,13,20,21,22,31,32]. Various studies have a different outcome for the combinations of drugs, like, in one study, a combination of sodium stibogluconate and plain amphotericin-B resulted in 2 deaths out of 4 cases (50% cure rate), while three drugs combination (sodium stibogluconate + plain amphotericin-B + miltefosine) for 33 cases resulted in all cure with one relapse which later treated with liposomal amphotericin-B (100% cure rate)[23].

For CL diagnosed cases use of intra-lesional sodium stibogluconate alone showed recovery of all 285 cases (100% cure rate)[8]. Inspiring results were also seen in cases where the combination of intravenous and intra-lesional stibogluconate resulted in the survival of all 18 cases[28].

In case of relapse or failure, liposomal or plain amphotericin-B was most commonly used, this showed diverse efficacy in different studies. Like in one, out of 10 cases, 6 survived, 3 Lost to follow up and 1 resulted in death after the use of plain amphotericin-B[26]. While in another study, plain

amphotericin-B was sufficient for the relapsed case after initial sodium stibogluconate (intralesional or intravenous not explained)[10]. A similar instance was reported in a study where plain amphotericin-B was given after failed miltefosine therapy and the case survived[10].

The dose of all drugs was not available in studies, however, a single dose (10 mg/kg) of liposomal amphotericin-B was used with a 100% cure rate including one relapse case after use of plain amphotericin-B[13].

DISCUSSION

The thirty-one studies of southern Himalayas show emerging leishmaniasis in high-altitude areas. The disease profile is distinctive from typical endemic areas. This can be discussed under various aspects of disease profile.

Epidemiology (demography)

Leishmaniasis is prevalent mainly in the poor and marginalized communities of the world, predominantly of the Indian subcontinent like Bangladesh, India, and Nepal. However, recent studies are suggestive of the emergence of new endemic foci in various parts of the world as well. In 2017, 94% of new VL cases were reported in seven countries: Brazil, Ethiopia, India, Kenya, Somalia, South Sudan, and Sudan while the majority of CL cases reported from Afghanistan, Algeria, Brazil, Colombia, the Islamic Republic of Iran, Pakistan, Peru, Saudi Arabia, and the Syrian Arab Republic[34]. Some latest sporadic cases have also been reported from Bhutan and Thailand[32]. All these countries share a similar topography, ecological and environmental factors (high humidity, adequate rainfall, and surface dampness) which are favorable for the proliferation of Phlebotomes. Results of recent studies demonstrate that now leishmaniasis is not confined to a specified topography, rainfall, temperature, or vegetation, it has now continuously expanded its geographical distribution which can be explained by factors such as rapid growing globalization, global warming, deforestation, and urbanization. These facts can't be confirmed as very few epidemiological studies are available on this issue. Furthermore, reviewing the literature, it was observed that the majority of the cases have been reported along with the major river belts in these new areas. This observation is highly suggestive of possible up stream migration of vectors along the rivers. In the past 15 years of reporting, good numbers of cases were found in newly endemic areas of Bhutan, Nepal, India (Uttarakhand, Himachal Pradesh, Jammu and Kashmir), and Northern parts of Pakistan.

Entomological and parasitological findings

Sandfly, vector of VL and CL, includes many species of the genus *Phlebotomus* (in the Old World) and *Lutzomyia longipalpis* (in the New World)[35,36]. Although the majority of the reviewed studies have not mentioned the associated vectors, *Phlebotomus argentipes* was found to be the predominant vector among the reported cases except in Himachal Pradesh (India) where *P. longiductus* and *P. major* were identified in co-existence. Interestingly *P. argentipes* remain closely associated with the exclusive cases of VL while *P. longiductus* (most common) and *P. major* were associated with areas where both CL & VL forms were found (Table 2). Therefore, the associated area needs an entomological study to know the basic characteristics of the vector and associated factors.

L. donovani transmission in East Africa consists of both anthroponotic and zoonotic components[37]. In Sudan, rodents and dogs were found to be reservoirs; however, observation in the majority of outbreaks reflects anthroponotic predominant transmission[38,39]. While in SEAR countries, the human being is the only reported reservoir. In this review also, we found a similar finding of the human being as the sole reservoir for VL.

Major species of parasites of VL are reported as *L. donovani* in South Asia and *L. infantum* in the Mediterranean region along with some sporadic cases in Central Asia, China, Mexico and Central Brazil [35,40]. The central western area of Brazil which is considered an area of recent transmission for VL and is on the risk for CL, *L. longipalpis* was the widespread species discovered[40]. In the new world, the most common etiological agent is *L. infantum*. The current review also documents similar findings of *L. donovani* in the majority of studies but one study from the Himalayan areas of Pakistan reported *L. infantum* in the majority[16]. A study in Brazil documented to have detected for the first time the presence of either *L. infantum* or *L. braziliensis* circulating in the domestic host[41]. In India, VL is caused by *L. donovani* in the north eastern region, and CL is caused by *L. tropica* in the western Thar Desert region[42]. Himachal Pradesh is a more recently leishmaniasis endemic state in north-west India where VL and CL coexist. The incidence of CL is higher than that of VL and most cases are attributable to *L. donovani*[33,43]. One of the studies conducted in the same region reported an interesting presence of *Leptomonas seymouri* co-infection in CL with *L. donovani*[30]. Undoubtedly there may be some missing links and associations that are still unknown and undiscovered since no other areas around Himachal Pradesh of the southern Himalayan region reported any remarkable epidemiological studies. Therefore, this review may act as a catalyst to perpetuate epidemiological search in this region to establish various niches.

Table 2 Characteristics of Leishmaniasis in the southern Himalayan region

Sr No.	Geographical area	Causative agent	Vector	Clinical picture
1	Northern areas of Pakistan[2]	<i>Leishmania infantum</i>	Not identified	Visceral leishmaniasis
2	Indian states of Jammu & Kashmir[3-5]	Not identified	Not identified	Cutaneous leishmaniasis most common with a single case study of visceral Leishmaniasis
3	Himachal Pradesh[6-16]	<i>L. donovani</i> & <i>L. tropica</i>	<i>P. longiductus</i> (most common) & <i>P. major</i>	Both cutaneous & visceral forms of Leishmaniasis
4	Uttarakhand (Garhwal)[17-21]	<i>L. donovani</i>	<i>P. argentipes</i>	Visceral leishmaniasis
5	Uttarakhand (Kumaon)[22-25]	Not identified	<i>P. argentipes</i>	Visceral leishmaniasis
6	Nepal[26-29]	<i>L. donovani</i>	<i>P. argentipes</i>	Visceral leishmaniasis
7	Bhutan[30,31]	<i>L. donovani</i>	<i>P. argentipes</i>	Visceral leishmaniasis

CL in the New World is generally caused by *L. mexicana*, while CL of the Old World is caused by five species of Leishmania: *L. infantum* (more common), *L. tropica*, *L. major*, *L. aethiopica*, and *L. donovani*. However, a study in the Indian sub-continent documents *L. tropica* in Pakistan, *L. donovani* and *L. major* in Nepal are the most common organism causing CL[4]. PKDL is caused primarily by *L. donovani* both in India and Sudan with only a few cases by *L. infantum* or *L. chagasi*[28].

This shows the existence of different types of species for both VL and CL in different parts of the South Asian countries including the southern Himalayas of the Indian Sub-continent. The rationale behind this diversity and associated epidemiological factors needs to be studied further.

Clinical presentation

VL has different clinical features in the endemic, epidemic, or sporadic situation. It tends to be relatively chronic and mostly affects children in endemic areas. Both the VL and CL are endemic in Pakistan and India while only VL is endemic in Nepal and Bhutan (WHO updates). Study analysis revealed that the characteristics of the disease vary with the environment. Here we see the preponderance of VL in Bhutan, Nepal, and Uttarakhand (India) with the coexistence of CL and VL in the Indian states of Himachal Pradesh, Jammu and Kashmir and Pakistan (Table 1).

Most cases are asymptomatic, but some eventually develop VL on follow-up, more commonly in males[35]. Risk factors for progression to VL include malnutrition, genetic factor and other co-infections, mainly HIV. The major classical presentation is prolonged fever, fatigue, loss of appetite and weight, and left hypochondrium discomfort. There may be non-tender splenomegaly with or without hepatomegaly, pallor, and lymphadenopathy (especially in Sudan, commonly by *Viannia* subgenus species). The darkening of the skin is typical for the Indian variant (Hindi name, kala-azar). Clinically CL usually exhibits painless, multiple, round-to-oval crater-form dry nodular lesions, mostly at the site of inoculation. Usually, these cutaneous lesions heal spontaneously in 1 year, often with disfiguring scars. PKDL is extremely rare, confined mainly in two regions endemic to kala-azar the Indian sub-continent and Sudan plus adjoining areas (up to 50% and 10% of patients with kala-azar respectively)[44-46].

Among all the studies reviewed none of them documented an asymptomatic period. The majority documented similar classical VL and CL symptoms except a few, which documented some atypical presentations like ascites, diarrhea, epistaxis, HLH syndrome, and hypergammaglobulinaemia (Table 1). Few cases of PKDL were reported from the hilly area of Uttarakhand too. The occurrence of PKDL after VL treatment in Nepal is also low as compared to neighboring countries[47].

Diagnosis

For diagnosis of Leishmaniasis many tests like dual path platform, a rapid immune-chromatographic test, and enzyme linked immune-sorbent assay (ELISA) are recommended by the Brazilian Ministry of Health[41]. Govt. of India recommends various tests for the detection of leishmania, including serology, aldehyde test, complement fixation test, indirect hem-agglutination test, ELISA, direct agglutination tests (DAT), spleen or bone marrow aspirates, and rK39[43]. The diagnostic policy for leishmaniasis is variable depending on the level of health systems. In first-line centers or rural hospitals of the highly endemic zone, the rK39 test is mostly used. Parasitological diagnosis is necessary for relapse identification. In low- endemic areas, more specific tests like PCR or parasitic demonstration are found necessary, as PCR is more sensitive than microscopic examination, therefore, can detect more asymptomatic infections. However, it is not available in most centers, and evaluation of its diagnostic accuracy and proper standardization is needed. For relapse, serological tests such as DAT, ELISA, and rK39 rapid test are usually positive and frequently used in majority areas but are of limited value, as a

positive result may be due to antibodies persisting after a past episode of VL, so better to show parasitological evidence for confirmation. A study in Brazil documented use of nested PCR (LnPCR) and PCR-restriction fragment length polymorphism for identification of *Leishmania* species[41]. A careful perusal of studies in this review showed a comprehensive use of various diagnostic procedures with no conclusive evidence towards any particular method. Future studies in these regions are need of the hour to formulate a diagnostic policy suitable for primary to tertiary health care levels.

Treatment and outcomes

Depending upon the sensitivity of drugs and the economic status, the treatment regime varies in different parts of the world. Liposomal amphotericin-B monotherapy (total dose of 20 to 21 mg/kg) is the preferred treatment in Europe, North America, and South America[48,49]. In East Africa, first-line therapy consists of a combination treatment of sodium stibogluconate and paromomycin for 17 d; the efficacy of liposomal amphotericin-B, miltefosine and paromomycin monotherapy are unacceptably low [50]. WHO Expert Committee and the Regional Technical Advisory Group of SEAR recommends liposomal amphotericin-B in a single dose of 10 mg/kg body weight as the first line treatment regimen for the Indian subcontinent within the current elimination strategy, given its high antimicrobial efficacy, safety, ease of use and assured compliance[1].

The majority of the studies in this review comply with the above standards and none of them documented parallel or supplemental pharmacotherapy other than the recommended regimen. However, agreeable documentation about the efficacy of the above drugs cannot be established in these emerging foci, as different studies had different outcomes. On summarizing the treatment outcomes liposomal amphotericin-B has emerged as the most effective therapy against the disease (with 100% cure rate achieved with single-dose). Furthermore, it is also found to be effective in VL-associated HLH and the explanations were that it inhibits macrophage function, reduces cytokine expression, and antigen-induced proliferation of T and B cells in vitro, causing a dual effect on both HLH and VL[13]. At last, the treatment regimen must follow national or regional guidelines, if applicable. Species identification usually is not critical to treatment decisions for VL (incontrast with CL)[51]. Multiple trial studies regarding drugs and doses should be done for the best suitable management protocol in these new niches.

Limitations

As said before, the availability of only a few studies related to the Himalayan regions is the major limitation of this review. Limited studies have covered the factors determining the transmission of VL in these new foci. The paucity of data limits the freedom to give any conclusive remarks on this new possible niche of leishmaniasis. A detailed analysis of these factors and the molecular characterization of vector species and leishmaniasis strain are still lacking. However, this mini-review aspires to highlight the surge of new cases in non-endemic areas as a matter of public health importance and research.

CONCLUSION

Despite substantial progress towards VL elimination in most endemic parts of the world, recently reported the emergence of new endemic foci in Southern Himalayas, forecast a great challenge for public health. Upstream river belts are a possible path of Sandfly spread towards these non-endemic areas, need a better environmental study to prove. In these areas, *P. argentipes* is found to be a predominant vector, *L. donovani* as a major parasite cause of VL, and *L. tropica*, *L. donovani*, and *L. major* as a major cause of CL in Pakistan, India, and Nepal respectively. Isolated VL is seen in Bhutan, Nepal, and the Uttarakhand state of India, while both VL and CL are seen in other Himalayan areas. Moreover, patients of these areas have a typical clinical presentations (ascites, diarrhea, epistaxis, HLH syndrome, and hypergammaglobulinaemia) so they need a high index of clinical suspicion, prompt diagnosis, and management. Single-dose liposomal amphotericin-B holds a 100% cure rate. As the a typical disease is recognized as a major threat to ongoing leishmaniasis elimination, so continuous monitoring of the disease type and associated parasitic variants and vector species should be implemented as part of the ongoing leishmaniasis elimination and maintenance programs. Studies on vector species and alternate reservoirs are also required for a better understanding of region-specific disease transmission and epidemiology.

FOOTNOTES

Author contributions: All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: India

ORCID number: Prasan Kumar Panda 0000-0002-3008-7245.

S-Editor: Gong ZM

L-Editor: A

P-Editor: Ju JL

REFERENCES

- 1 **World Health Organization.** Kala-Azar elimination program: report of a WHO consultation of partners, Geneva, Switzerland, February 10-11, 2015. Available from: https://apps.who.int/iris/bitstream/handle/10665/185042/9789241509497_eng.pdf?sequence=1
- 2 **Hirve S, Kroeger A, Matlashewski G, Mondal D, Banjara MR, Das P, Be-Nazir A, Arana B, Oliaro P.** Towards elimination of visceral leishmaniasis in the Indian subcontinent-Translating research to practice to public health. *PLoS Negl Trop Dis* 2017; **11**: e0005889 [PMID: 29023446 DOI: 10.1371/journal.pntd.0005889]
- 3 **Raina S, Mahesh DM, Kaul R, Satindera KS, Gupta D, Sharma A, Thakur S.** A new focus of visceral leishmaniasis in the Himalayas, India. *J Vector Borne Dis* 2009; **46**: 303-306 [PMID: 19959858]
- 4 **Katakura K.** Molecular epidemiology of leishmaniasis in Asia (focus on cutaneous infections). *Curr Opin Infect Dis* 2009; **22**: 126-130 [PMID: 19276879 DOI: 10.1097/QCO.0b013e3283229ff2]
- 5 **Verma SK, Ahmad S, Shirazi N, Kusum A, Kaushik RM, Barthwal SP.** Sodium stibogluconate-sensitive visceral leishmaniasis in the non-endemic hilly region of Uttarakhand, India. *Trans R Soc Trop Med Hyg* 2007; **101**: 730-732 [PMID: 17382981 DOI: 10.1016/j.trstmh.2007.02.008]
- 6 **Rao JS, Sharma SK, Bhattacharya D, Saxena NB.** Sandfly survey in Nainital and Almora districts of Uttaranchal with particular reference to *Phlebotomus argentipes*, vector of kala-azar. *J Commun Dis* 2001; **33**: 7-11 [PMID: 11898464]
- 7 **Bhattacharya SK, Rinzin N, Chusak P, Dash AP, Chowdhury R, Tobgay T, Narain JP.** Occurrence & significance of kala-azar in Bhutan. *Indian J Med Res* 2010; **132**: 337-338 [PMID: 20847382]
- 8 **Sharma NL, Mahajan VK, Negi AK.** Epidemiology of a new focus of localized cutaneous leishmaniasis in Himachal Pradesh. *J Commun Dis* 2005; **37**: 275-279 [PMID: 17278657]
- 9 **Sharma NL, Mahajan VK, Kanga A, Sood A, Katoch VM, Mauricio I, Singh CD, Parwan UC, Sharma VK, Sharma RC.** Localized cutaneous leishmaniasis due to *Leishmania donovani* and *Leishmania tropica*: preliminary findings of the study of 161 new cases from a new endemic focus in himachal pradesh, India. *Am J Trop Med Hyg* 2005; **72**: 819-824 [PMID: 15964970]
- 10 **Yangzom T, Cruz I, Bern C, Argaw D, den Boer M, Vélez ID, Bhattacharya SK, Molina R, Alvar J.** Endemic transmission of visceral leishmaniasis in Bhutan. *Am J Trop Med Hyg* 2012; **87**: 1028-1037 [PMID: 23091191 DOI: 10.4269/ajtmh.2012.12-0211]
- 11 **Thakur L, Singh KK, Kushwaha HR, Sharma SK, Shankar V, Negi A, Verma G, Kumari S, Jain A, Jain M.** *Leishmania donovani* Infection with Atypical Cutaneous Manifestations, Himachal Pradesh, India, 2014-2018. *Emerg Infect Dis* 2020; **26**: 1864-1869 [PMID: 32687048 DOI: 10.3201/eid2608.191761]
- 12 **Sharma RC, Mahajan VK, Sharma NL, Sharma A.** A new focus of cutaneous leishmaniasis in Himachal Pradesh (India). *Indian J Dermatol Venereol Leprol* 2003; **69**: 170-172 [PMID: 17642870]
- 13 **Kumari S, Dhawan P, Panda PK, Bairwa M, Pai VS.** Rising visceral leishmaniasis in Holy Himalayas (Uttarakhand, India) - A cross-sectional hospital-based study. *J Family Med Prim Care* 2020; **9**: 1362-1369 [PMID: 32509616 DOI: 10.4103/jfmpe.jfmpe_1174_19]
- 14 **Ostyn B, Uranw S, Bhattarai NR, Das ML, Rai K, Tersago K, Pokhrel Y, Durnez L, Marasini B, Van der Auwera G, Dujardin JC, Coosemans M, Argaw D, Boelaert M, Rijal S.** Transmission of *Leishmania donovani* in the Hills of Eastern Nepal, an Outbreak Investigation in Okhaldhunga and Bhojpur Districts. *PLoS Negl Trop Dis* 2015; **9**: e0003966 [PMID: 26252494 DOI: 10.1371/journal.pntd.0003966]
- 15 **Pandey BD, Pun SB, Kaneko O, Pandey K, Hirayama K.** Case report: Expansion of visceral leishmaniasis to the western hilly part of Nepal. *Am J Trop Med Hyg* 2011; **84**: 107-108 [PMID: 21212211 DOI: 10.4269/ajtmh.2011.10-0291]
- 16 **Rab MA, Evans DA.** *Leishmania infantum* in the Himalayas. *Trans R Soc Trop Med Hyg* 1995; **89**: 27-32 [PMID: 7747300 DOI: 10.1016/0035-9203(95)90644-4]
- 17 **Sharma NL, Mahajan VK, Negi AK, Verma GK.** The rK39immunochromatic dipstick testing: a study for K39 seroprevalence in dogs and human leishmaniasis patients for possible animal reservoir of cutaneous and visceral leishmaniasis in endemic focus of Satluj river valley of Himachal Pradesh (India). *Indian J Dermatol Venereol Leprol* 2009; **75**: 52-55 [PMID: 19172032 DOI: 10.4103/0378-6323.45221]
- 18 **Mahajan D, Bhat ML, Singh JB, Hans D.** Visceral Leishmaniasis In A Native Kashmiri Boy. *JK Sci* 2009; **11**: 152-153

- 19 **Raina S**, Raina RK, Sharma R, Rana BS, Bodh A, Sharma M. Expansion of visceral leishmaniasis to northwest sub-Himalayan region of India: A case series. *J Vector Borne Dis* 2016; **53**: 188-191 [PMID: 27353591]
- 20 **Raina RK**, Raina S, Sharma M. Visceral leishmaniasis-associated hemophagocytosis: A tale of two unexpected diagnoses from a nonendemic region. *Trop Parasitol* 2017; **7**: 56-58 [PMID: 28459018 DOI: 10.4103/2229-5070.202288]
- 21 **Pawar S**, Ragesh R, Nischal N, Sharma S, Panda PK, Sharma SK. Unique Triad of 'Pregnancy, Kala Azar and Hemophagocytic Lymphohistiocytic Syndrome from a Non-Endemic Region'. *J Assoc Physicians India* 2015; **63**: 65-68 [PMID: 26710404]
- 22 **Chandra H**, Chandra S, Kaushik R, Bhat N, Shrivastava V. Hemophagocytosis on bone marrow aspirate cytology: single center experience in north himalayan region of India. *Ann Med Health Sci Res* 2014; **4**: 692-696 [PMID: 25328776 DOI: 10.4103/2141-9248.141515]
- 23 **Ahmad S**, Chandra H, Bhat NK, Dhar M, Shirazi N, Verma SK. North Indian state of Uttarakhand: a new hothouse of visceral leishmaniasis. *Trop Doct* 2016; **46**: 111-113 [PMID: 26466848 DOI: 10.1177/0049475515609245]
- 24 **Kumar Bhat N**, Ahuja V, Dhar M, Ahmad S, Pandita N, Gupta V, Chandra S. Changing Epidemiology: A New Focus of Kala-azar at High-Altitude Garhwal Region of North India. *J Trop Pediatr* 2017; **63**: 104-108 [PMID: 27582128 DOI: 10.1093/tropej/fmw056]
- 25 **Singh S**, Biswas A, Wig N, Aggarwal P, Sood R, Wali JP. A new focus of visceral leishmaniasis in sub-Himalayan (Kumaon) region of northern India. *J Commun Dis* 1999; **31**: 73-77 [PMID: 10810593]
- 26 **Kumar A**, Rawat V, Thapliyal N, Saxena SR. Kala-azar-A case series from the nonendemic area, Uttarakhand. *Ann Trop Med Public Health* 2013; **6**: 355-357 [DOI: 10.4103/1755-6783.121008]
- 27 **Mathur SB**, Arya AK. Nonmigrant children with visceral leishmaniasis from the nonendemic area of Uttarakhand. *J Trop Pediatr* 2014; **60**: 322-325 [PMID: 24531375 DOI: 10.1093/tropej/fmu007]
- 28 **Wani GM**, Ahmad SM, Khursheed B. Clinical study of cutaneous leishmaniasis in the Kashmir Valley. *Indian Dermatol Online J* 2015; **6**: 387-392 [PMID: 26753136 DOI: 10.4103/2229-5178.169732]
- 29 **Leherwal MA**, Yasin SB, Ahmed SB. Diagnosis of cutaneous leishmaniasis by FNAC-Report of three cases. *J Cytol* 2004; **21**: 103-105
- 30 **Thakur L**, Kushwaha HR, Negi A, Jain A, Jain M. Leptomonas seymouri Co-infection in Cutaneous Leishmaniasis Cases Caused by Leishmania donovani From Himachal Pradesh, India. *Front Cell Infect Microbiol* 2020; **10**: 345 [PMID: 32760679 DOI: 10.3389/fcimb.2020.00345]
- 31 **Pandey BD**, Pandey K, Kaneko O, Yanagi T, Hirayama K. Relapse of visceral leishmaniasis after miltefosine treatment in a Nepalese patient. *Am J Trop Med Hyg* 2009; **80**: 580-582 [PMID: 19346379]
- 32 **Schwarz D**, Andrews J, Gauchan B. Visceral leishmaniasis in far western Nepal: another case and concerns about a new area of endemicity. *Am J Trop Med Hyg* 2011; **84**: 508 [PMID: 21363996 DOI: 10.4269/ajtmh.2011.11-0021]
- 33 **Sharma NL**, Sood A, Arora S, Kanga A, Mahajan V, Negi AK, Sharma AK. Characteristics of Leishmania spp. isolated from a mixed focus of cutaneous and visceral leishmaniasis in Himachal Pradesh (India). *Int J Third World Med* 2009; **7** [DOI: 10.5580/1683]
- 34 **World Health Organization**. Visceral leishmaniasis-WHO publishes validation documents as countries approach elimination. Available from: https://www.who.int/neglected_diseases/news/Visceral_leishmaniasis_WHO_publishes_validation_document/en/
- 35 **World Health Organization**. Control of the Leishmaniasis: Report of the WHO Expert Committee Meeting, Geneva. March 22-26, 2010. Available from: <https://apps.who.int/iris/handle/10665/44412>
- 36 **Killick-Kendrick R**. Phlebotomine vectors of the leishmaniasis: a review. *Med Vet Entomol* 1990; **4**: 1-24 [PMID: 2132963 DOI: 10.1111/j.1365-2915.1990.tb00255]
- 37 **Alvar J**, Bashaye S, Argaw D, Cruz I, Aparicio P, Kassa A, Orfanos G, Parreño F, Babaniyi O, Gudeta N, Cañavate C, Bern C. Kala-azar outbreak in Libo Kemkem, Ethiopia: epidemiologic and parasitologic assessment. *Am J Trop Med Hyg* 2007; **77**: 275-282 [PMID: 17690399 DOI: 10.4269/ajtmh.2007.77.275]
- 38 **Bucheton B**, Kheir MM, El-Safi SH, Hammad A, Mergani A, Mary C, Abel L, Dessein A. The interplay between environmental and host factors during an outbreak of visceral leishmaniasis in eastern Sudan. *Microbes Infect* 2002; **4**: 1449-1457 [PMID: 12475635 DOI: 10.1016/S1286-4579(02)00027-8]
- 39 **Dereure J**, El-Safi SH, Bucheton B, Boni M, Kheir MM, Davoust B, Pratlong F, Feugier E, Lambert M, Dessein A, Dedet JP. Visceral leishmaniasis in eastern Sudan: parasite identification in humans and dogs; host-parasite relationships. *Microbes Infect* 2003; **5**: 1103-1108 [PMID: 14554251 DOI: 10.1016/j.micinf.2003.07.003]
- 40 **Bern C**, Maguire JH, Alvar J. Complexities of assessing the disease burden attributable to leishmaniasis. *PLoS Negl Trop Dis* 2008; **2**: e313 [PMID: 18958165 DOI: 10.1371/journal.pntd.0000313]
- 41 **Menezes JA**, Ferreira Ede C, Andrade-Filho JD, de Sousa AM, Morais MH, Rocha AM, Machado-Coelho GL, Lima FP, Madureira AP, Garcia TC, Freitas CR, Soares RP, Margonari C. An Integrated Approach Using Spatial Analysis to Study the Risk Factors for Leishmaniasis in Area of Recent Transmission. *Biomed Res Int* 2015; **2015**: 621854 [PMID: 26229961 DOI: 10.1155/2015/621854]
- 42 **Thakur L**, Singh KK, Shanker V, Negi A, Jain A, Matlashewski G, Jain M. Atypical leishmaniasis: A global perspective with emphasis on the Indian subcontinent. *PLoS Negl Trop Dis* 2018; **12**: e0006659 [PMID: 30260957 DOI: 10.1371/journal.pntd.0006659]
- 43 **National Center for Vector Borne Diseases Control (NCVBDC)**. Guidelines-Diagnosis and treatment of Kala-azar. [Accessed on 10 October 2021] Available from: <http://nvbdcp.gov.in/Doc/Guidelines-Diagnosis-Treatment-KA.pdf>
- 44 **Baghestani S**, Handjani F, Sodeifi M, Kumar PV. Post-kala-azar dermal leishmaniasis. *Eur J Dermatol* 1998; **8**: 277-279 [PMID: 9649686]
- 45 **Croft SL**. PKDL--a drug related phenomenon? *Indian J Med Res* 2008; **128**: 10-11 [PMID: 18820352]
- 46 **Zijlstra EE**, Musa AM, Khalil EA, el-Hassan IM, el-Hassan AM. Post-kala-azar dermal leishmaniasis. *Lancet Infect Dis* 2003; **3**: 87-98 [PMID: 12560194 DOI: 10.1016/S1473-3099(03)00517-6]
- 47 **Uranw S**, Ostyn B, Rijal A, Devkota S, Khanal B, Menten J, Boelaert M, Rijal S. Post-kala-azar dermal leishmaniasis in

- Nepal: a retrospective cohort study (2000-2010). *PLoS Negl Trop Dis* 2011; **5**: e1433 [PMID: [22206030](#) DOI: [10.1371/journal.pntd.0001433](#)]
- 48 **Aronson N**, Herwaldt BL, Libman M, Pearson R, Lopez-Velez R, Weina P, Carvalho E, Ephros M, Jeronimo S, Magill A. Diagnosis and Treatment of Leishmaniasis: Clinical Practice Guidelines by the Infectious Diseases Society of America (IDSA) and the American Society of Tropical Medicine and Hygiene (ASTMH). *Am J Trop Med Hyg* 2017; **96**: 24-45 [PMID: [27927991](#) DOI: [10.4269/ajtmh.16-84256](#)]
- 49 **Alvar J**, Croft S, Olliaro P. Chemotherapy in the treatment and control of leishmaniasis. *Adv Parasitol* 2006; **61**: 223-274 [PMID: [16735166](#) DOI: [10.1016/S0065-308X\(05\)61006-8](#)]
- 50 **Musa A**, Khalil E, Hailu A, Olobo J, Balasegaram M, Omollo R, Edwards T, Rashid J, Mbui J, Musa B, Abuzaid AA, Ahmed O, Fadlalla A, El-Hassan A, Mueller M, Mucee G, Njoroge S, Manduku V, Mutuma G, Apadet L, Lodenyo H, Mutea D, Kirigi G, Yifru S, Mengistu G, Hurissa Z, Hailu W, Weldegebreal T, Tafes H, Mekonnen Y, Makonnen E, Ndegwa S, Sagaki P, Kimutai R, Kesusu J, Owiti R, Ellis S, Wasunna M. Sodium stibogluconate (SSG) & paromomycin combination compared to SSG for visceral leishmaniasis in East Africa: a randomised controlled trial. *PLoS Negl Trop Dis* 2012; **6**: e1674 [PMID: [22724029](#) DOI: [10.1371/journal.pntd.0001674](#)]
- 51 **Sundar S**, Singh A. Recent developments and future prospects in the treatment of visceral leishmaniasis. *Ther Adv Infect Dis* 2016; **3**: 98-109 [PMID: [27536354](#) DOI: [10.1177/2049936116646063](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

