

Format for Manuscript Preparation and Submission: Meta-Analysis

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO:

Manuscript Type: META-ANALYSIS

Effects of neoadjuvant chemotherapy *vs* chemoradiotherapy in the treatment of esophageal adenocarcinoma: A systematic review and meta-analysis

Csontos A *et al.* Impact of neoadjuvant therapy of esophageal adenocarcinoma

Armand Csontos, Alíz Fazekas, Lajos Szakó, Nelli Farkas, Csenge Papp, Szilárd Ferenczi, Szabolcs Bellyei, Péter Hegyi, András Papp

Armand Csontos, Csenge Papp, Szilárd Ferenczi, András Papp, Department of Surgery, University of Pécs, Medical School, Clinical Center, Pécs H-7624, Baranya, Hungary

Alíz Fazekas, Nelli Farkas, Institute of Bioanalysis, University of Pécs, Medical School, Pécs H-7624, Baranya, Hungary

Alíz Fazekas, Nelli Farkas, Péter Hegyi, Institute for Translational Medicine, University of Pécs, Medical School, Pécs H-7624, Baranya, Hungary

Lajos Szakó, Department of Emergency Medicine, Clinical Center, University of Pécs, Medical School, Pécs 7624, Baranya, Hungary

Szabolcs Bellyei, Department of Oncotherapy, University of Pécs, Medical School, Clinical Center, Pécs H-7624, Baranya, Hungary

Péter Hegyi, Centre for Translational Medicine, Semmelweis University, Budapest 1085, Hungary

Péter Hegyi, Institute of Pancreatic Diseases, Semmelweis University, Budapest H-1083, Hungary

Co-first authors:

Co-corresponding authors:

Author contributions: Csontos A contributed to the design and implementation of the study and the writing of the manuscript; Fazekas A and Farkas N contributed to the statistical analyses and the writing of the manuscript; Szakó L contributed to the design of the study and the revision of the manuscript; Papp C and Ferenczi S contributed to the performance of the research; Bellyei S and Hegyi P contributed to the quality and professional revision; Papp A contributed to the quality and professional revision and the writing of the manuscript.

Supported by

Corresponding author: Armand Csontos, MD, Doctor, Department of Surgery, University of Pécs, Medical School, Clinical Center, Ifjúság Street 13, Pécs H-7624, Baranya, Hungary. csontos.armand@gmail.com

Received: December 23, 2023

Revised:

Accepted:

Published online:

Abstract

BACKGROUND

Neoadjuvant therapy is an essential modality for reducing the clinical stage of esophageal cancer; however, the superiority of neoadjuvant chemotherapy (nCT) or neoadjuvant chemoradiotherapy (nCRT) is unclear. Therefore, a discussion of these two modalities is necessary.

AIM

To investigate the benefits and complications of neoadjuvant modalities.

METHODS

To address this concern, predefined criteria were established using the PICO protocol. Two independent authors performed comprehensive searches using predetermined keywords. Statistical analyses were performed to identify significant differences between groups. Potential publication bias was visualized using funnel plots. The quality of the data was evaluated using the Risk of Bias Tool 2 (RoB2) and the GRADE approach.

RESULTS

Ten articles, including 1928 patients, were included for the analysis. Significant difference was detected in pathological complete response (pCR) [$P < 0.001$; odds ratio (OR): 0.27; 95%CI: 0.16-0.46], 30-d mortality ($P = 0.015$; OR: 0.4; 95%CI: 0.22-0.71) favoring the nCRT, and renal failure ($P = 0.039$; OR: 1.04; 95%CI: 0.66-1.64) favoring the nCT. No significant differences were observed in terms of survival, local or distal recurrence, or other clinical or surgical complications. The result of RoB2 was moderate, and that of the GRADE approach was low or very low in almost all cases.

CONCLUSION

Although nCRT may have a higher pCR rate, it does not translate to greater long-term survival. Moreover, nCRT is associated with higher 30-d mortality, although the specific cause for postoperative complications could not be identified. In the case of nCT, toxic side effects are suspected, which can reduce the quality of life. Given the quality of available studies, further randomized trials are required.

Key Words: Neoadjuvant; Chemotherapy; Chemoradiotherapy; Esophageal cancer; Adenocarcinoma

Csontos A, Fazekas A, Szakó L, Farkas N, Papp C, Ferenczi S, Bellyei S, Hegyi P, Papp A. Effects of neoadjuvant chemotherapy vs chemoradiotherapy in the treatment of esophageal adenocarcinoma: A systematic review and meta-analysis. *World J Gastroenterol* 2024; 0(0): 0000-0000

URL: <https://www.wjgnet.com/1007-9327/full/v0/i0/0000.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v0.i0.0000>

Core Tip: Neoadjuvant chemoradiation increases pathological complete response and 30-d mortality in patients with esophageal adenocarcinoma; however, it has no effect on long-term survival. It may be associated with side effects that can reduce the quality of life.

INTRODUCTION

Epidemiology

Esophageal cancer (EC) is the eighth most prevalent cancer, with more than 500000 cases worldwide, and it is the sixth leading cause of tumor mortality. Squamous cell carcinoma (SCC) is still the leading subtype in the Asian EC Belt; however, in Western countries,

such as North America, Oceania, and Western and Northern Europe, including Hungary, the incidence rate of adenocarcinoma (AC) has been increasing, surpassing that of SCC[1,2].

In the early stages, surgery can lead to full recovery; however, an advanced tumor stage at initial diagnosis can result in high morbidity and mortality rates[3]. Esophagectomy with radical lymphadenectomy is one of the most invasive gastrointestinal procedures. To improve treatment results, a multidisciplinary approach is important, including the application of the enhanced recovery after surgery protocol[4,5], the minimally invasive approach of esophagectomy[6], and neoadjuvant oncological therapy, which can decrease mortality by 25%-35% compared with that of surgery alone[7-9].

Impact of the topic

The superiority of neoadjuvant therapy has been proven in several meta-analyses[7-9]. Neoadjuvant chemotherapy (nCT) or preoperative neoadjuvant chemoradiotherapy (nCRT) can also improve oncologic endpoints[8-15], increase overall and progression-free surveillance, and pathological complete response (pCR); however, it may also be associated with numerous clinical or surgical side effects and impaired quality of life. Therefore, the cost-benefit balance of these modalities is still unclear, especially in cases of AC of the esophagus and esophagogastric junction (GEJ).

Literature background

Previous meta-analyses have numerous limitations, including patients with SCC and AC as a homogenous population. Therefore, the results cannot be clearly applied to either subtype.

Impact of our analysis

We performed a comprehensive, up-to-date investigation to determine whether nCT or nCRT yields more favorable results in the surgical treatment of AC of the esophagus and esophageal junction.

MATERIALS AND METHODS

Protocol registration

The objectives and methodologies of this meta-analysis were predefined in a protocol registered with PROSPERO[16]. The registration was accepted on November 01, 2023, under the number CRD42023478615.

Question of the review

To define the scope of this meta-analysis, we used the PICO protocol, focusing on patients with esophageal or cardiac AC, who received neoadjuvant therapy before surgery. Intervention assessed was preoperative nCT, which was compared to nCRT. We investigated the following outcomes: Survival, remission rate, mortality, short- and long-term clinical and surgical complications, and quality of life. First, we planned to investigate only randomized controlled trials (RCTs) to minimize the risk of bias; however, to achieve an adequate sample size and robust conclusions, propensity score matched and high-quality cohort studies were also included. Studies that did not strictly involve patients with AC were excluded.

Search strategy and search terms

We conducted a comprehensive search on September 15, 2023, using PubMed, Embase, Cochrane, Web of Science, and Scopus databases. We used previously defined search terms, including “neoadjuvant,” “chemotherapy,” “chemoradiotherapy,” “esophageal cancer,” “esophagectomy,” and other random keywords, and their variants. The retrieved datasets were imported into the EndNote (ver. x9.3.3; Alfasoft AB, Göteborg, Sweden) library.

Selection process

Two independent authors conducted the selection process using EndNote software. The Cohen's kappa coefficients were calculated from these results. Discrepancies were resolved through consensus.

Data extraction

Data were extracted from text, figures, and tables of the included articles by two independent authors, with any discrepancies resolved through mutual agreement. Plot digitizer applications were used to collect data not provided in a numerical format[17]. Excel (Office 365, Microsoft, Redmond, WA, United States) datasheets were used to collect and organize the datasets. Descriptive data collected included study characteristics (author, year, type, and number of elements), patients demographics (age, sex, and performance), tumors (stage, location), and therapy (neoadjuvant regimen, surgical procedure). A meta-analysis was performed on outcomes with at least four homogeneous datasets. Outcomes, ineligible for statistical analysis, were qualitatively described. The outcomes assessed included pCR, surveillance (overall, progression-free, disease-free), mortality (30 or 90 d), tumor remission (local or distant), clinical complications (thromboembolism, respiratory and cardiac complications, renal failure, neutropenia) and surgical complication (anastomotic and chyle leakage, wound infection, bleeding, vocal cord paresis).

Statistical analysis

A random-effects meta-analysis was performed. Odds ratios (OR) with 95%CI were calculated to measure the effect size. To calculate the OR and pooled odds ratio, data for the total number of patients and those experiencing the event of interest in each group separately (referred as "raw data") was extracted or calculated from the studies, where it was available. The results are presented as the odds of an event of interest in the experimental group *vs* the control group. The results were considered statistically significant if the pooled CI did not contain a null value. We also performed a

supplementary analysis. Using the WebPlotDigitizer online tool, we digitalized the Kaplan-Meier (KM) curves published in the involved studies. Then, by applying the methodology of Guyot *et al*[18,19], we estimated the individual patient time-to-event data. Finally, we plotted all the available KM curves in the same figure. Using the estimated raw data, we calculated the hazard ratio (HR) within the studies and the pooled HR. A less than one HR suggests a smaller risk in the experimental group. The HR result was considered significant if it was not included in the confidence interval.

We visualized the findings in forest plots. Where applicable-the study number was large enough and not too heterogeneous-we also reported the prediction intervals (*i.e.*, the expected range of effects of future studies) of the results. Additionally, between-study heterogeneity was described using Higgins and Thompson's (*I*²) statistics (Higgins and Thompson[20], in 2002).

Publication bias was assessed by visual inspection of the funnel plots and calculation of the Harbord (modified Egger's) test *P* value[21] for the OR effect size. We assumed the presence of a possible small study bias if the *P* value was < 10%. However, we kept in mind that the test has limited diagnostic assessment (below 10 studies). Potential outlier publications were explored using different influence measures and plots following the recommendations of Harrer *et al*[22]. All statistical analyses were performed with R (R Core Team 2023, v4.3.0)[23] using the meta (Schwarzer 2023, v6.2.1)[24] package for basic meta-analysis calculations and plots, IPDfromKM for raw data simulations, and the dmetar (v0.0.9000)[25] package for additional influential analysis calculations and plots. To pool the effect size, the pooled OR based on raw data was calculated using the Mantel-Haenszel method[26,27]. The Exact Mantel-Haenszel method (without continuity correction) was used to handle zero-cell counts[28,29]. We used the Hartung-Knapp adjustment[30,31] for the CIs. To estimate the heterogeneity variance measure for the raw data OR calculation, the Paule-Mandel method[32] (recommended by Veroniki *et al*[33]) was used with the Q profile method for the confidence interval. Prediction interval calculations were based on the t-distribution. In the case of 0 cell counts, individual study OR with 95%CI was calculated by adding 0.5, as continuity correction (it was used only

for visualization on forest plot). The pooled HR was calculated using classical inverse-variance meta-analysis of log-transformed HR ratios using the REML heterogeneity variance estimator.

Descriptive analyses were performed by calculating the means, standard deviations, and percentages. The mean estimates from the median and range were calculated as follows[34]:

Risk of bias and certain of evidence

The Risk of Bias Tool 2 (RoB2) and GRADE approaches were used to assess the quality of the articles and our research.

RESULTS

Search process

A total of 1285 articles were identified from the five databases. After removing duplicates, 1141 articles were screened, after which 485 and 153 articles were selected based on title and abstract screening, respectively. Subsequently, 125 full-text reports were examined, and eight studies were included in the quantitative synthesis. Cohen's kappa indicated 99.74% substantial agreement (Cohen's k : 0.77). Some reports could not be retrieved as they were conference abstracts[28]. Articles were excluded based on predefined criteria (83 articles), including those covering only SCCs (9 articles) or mixed group of ACs and SCCs (23 articles), mentioning no pathological subtype (1 article), and being a preliminary trial (1 article). Two additional articles were included during the screening of previous reviews. Overall, the analysis included ten articles. More information is provided in Figure 1.

Characteristics of the studies

Ten articles, published between 2011 and 2018, were included in this meta-analysis. Seven studies were conducted in Europe, two in Australia, and one in the United States[35-44]. Of the 10 studies, two were RCTs[43,44], four were propensity score-matched cohort

study[39-42], and four were cohort study based on prospective institutional databases (clinical cancer registry)[35-38]. Six studies were single-center trials[36,39,40,42-44]. The articles collectively included data from 1928 patients, with 956 and 972 patients the nCT and nCRT groups, respectively. All included patients had esophageal ACs. Additional details are provided in Table 1.

Characteristics of the patients

The estimated mean age of the patients in both the groups was 60 years. The age range was 12-84 and 19-83 years. The nCT and nCRT groups included 829 (91%) and 857 (94%) male patients, respectively. Based on the available data, 84%, 16% and < 0.1% of the patients had American Society of Anesthesiologists scores of I-II, III, and IV, respectively. The patients had coronary morbidity (18%), diabetes mellitus (16%), pulmonary morbidity, chronic obstructive pulmonary disease (9%), history of malignancy (6%), and history of smoking (42%). More detailed information is summarized in Supplementary Table 1.

Characteristics of the tumor and pathological approach

Based on the available data, 99% of the tumors were diagnosed in the lower third of the esophagus or the GEJ. Clinical T-stages 1-4 accounted for 1%, 16%, 80%, and 3% of the cases, respectively. Nodal involvement was observed in 367 (61%) and 350 patients (59%) in the nCT and nCRT groups, respectively. Tumor differentiation was good in 2% and 1%, moderate in 36% and 31%, and poor in 57% and 64% of patients in the nCT and nCRT groups, respectively. Margin negative resection (R0) was performed in 696 (81%) and 800 (92%) patients in the nCT and nCRT groups. Pathological T-stage 1-4 accounted for 13%, 15%, 22%, and 47% of the cases, respectively, whereas N-stages 0-3 accounted for 45%, 31%, 14%, and 9% of the cases, respectively. Tumor regression grade (TRG, Mandard) stages 1-4/5 (in the nCT and nCRT groups) accounted for 14% (6%-22%), 17% (7%-26%), 24% (18%-29%), and 42% (62%-22%) of the cases, respectively additional details are provided in Supplementary Table 2.

Characteristics of the neoadjuvant therapy

Neoadjuvant regimens were administered to patients in both groups. The most frequently used neoadjuvant drugs were cisplatin, 5-fluorouracil, and docetaxel. The CROSS protocol was the most commonly used protocol in the chemoradiation group. Additional details are provided in Table 2[7,8,10,45-48].

Characteristics of the surgical procedure

Based on the available data, Ivor-Lewis (transthoracic), Orringer (transhiatal), McKeown (thoraco-abdomino-cervical) esophagectomies, and total gastrectomy were performed in 67%, 23%, 5%, and 4% of the patients, respectively. Minimally invasive or hybrid surgery techniques were performed in 27% and 51% of the patients, respectively, and open surgery was performed in only 23% of the patients. Two-field lymphadenectomy was the standard procedure in 74% of the patients, whereas three-field lymphadenectomy was performed in only 5% of the patients. Additional details are provided in Supplementary Table 3.

Pathological complete response

Data from eight studies, covering a total of 1547 patients, were analyzed[35,36,39-44]. The OR (pooled effect size) was 0.27 (95%CI: 0.16-0.46). A significant difference was observed, favoring nCRT over nCT ($P < 0.001$). Between-study heterogeneity, expressed as the I^2 value, was 0.29 (95%CI: 0-0.68; Figure 2).

Thirty-day mortality

Data from four studies, including 899 patients, were analyzed[38,40,41,43]. The OR was 0.4 (95%CI: 0.22-0.71). A significant difference was observed, favoring nCRT over nCT ($P = 0.015$), with no between-study heterogeneity (I^2 value: 0; 95%CI: 0-0.85; Figure 3).

Ninety-day mortality

Data from four studies, encompassing 108 patients, were analyzed[38,40-42]. The OR was 0.71 (95%CI: 0.28-1.84). No significant difference was observed between nCRT and nCT ($P = 0.34$), with no between-study heterogeneity (I^2 value: 0; 95%CI: 0-0.85; Supplementary Figure 1).

Overall survival

The KM curves and logHR analysis conducted for eight studies encompassing 1540 patients did not reveal significant differences between the two groups in terms of overall survival ($P = 0.82$)[36-42,44]. The OR was 0.98 (95%CI: 0.77-1.23) and between-study heterogeneity was 0.35 (95%CI: 0-0.71; Figures 4 and 5).

Considering the 12-month overall survival (OS), nine studies including 1588 patients were selected for analysis[36-44]. The OR was 1.08 (95%CI: 0.8-1.46). No significant difference was observed between the two groups ($P = 0.551$). The between-study heterogeneity was 0.05 (95%CI, 0-0.67; Supplementary Figure 2). The log HR analysis revealed no significant differences between the groups (Supplementary Figure 3).

For the 24-month OS, the OR was 1.03 (95%CI: 0.73-1.45)[36-44]. No significant difference was observed between the two groups ($P = 0.858$). The between-study heterogeneity was 0.42 (95%CI: 0-0.73; Supplementary Figure 4). The log HR analysis revealed no significant differences between the groups (Supplementary Figure 5).

Considering the 36-month OS, the OR was 0.93 (95%CI: 0.54-1.6)[36-44]. No significant difference was observed between the two groups ($P = 0.754$). The between-study heterogeneity was 0.73 (95%CI: 0.47-0.86; Supplementary Figure 6). The logHR analysis revealed no significant differences between the groups (Supplementary Figure 7).

Considering the 48-month OS, seven studies including 1066 patients were selected for analysis[36-38,40,42-44]. The OR was 0.67 (95%CI: 0.27-0.85). No significant difference was observed between the two groups ($P = 0.616$). The between-study heterogeneity was 0.67 (95%CI: 0.27-0.85; Supplementary Figure 8). The log HR analysis revealed no significant differences between the groups (Supplementary Figure 9).

Considering the 60-month OS, the OR was 1.15 (95%CI: 0.56-2.35)[36,38,40,42-44]. No significant difference was observed between the two groups ($P = 0.658$). The between-study heterogeneity was 0.67 (95%CI: 0.27-0.85; Supplementary Figure 10).

Disease-free survival

The KM curves and logHR analysis conducted for two studies including 578 patients did not reveal significant differences in overall survival between the two groups ($P = 0.85$)[36,38,42]. The OR was 1.04 (95%CI: 0.5-2.16). The between-study heterogeneity was 0.49 (95%CI: 0-0.85; Supplementary Figures 11 and 12).

Considering the 12-month Disease-free survival (DFS), the OR was 0.93 (95%CI: 0.44-1.97)[36,38,42]. No significant difference was observed between the two groups ($P = 0.702$). The between-study heterogeneity was 0.07 (95%CI: 0-0.9; Supplementary Figure 13). The logHR analysis revealed no significant differences between the groups (Supplementary Figure 14).

Considering the 24-month DFS, the OR was 0.95 (95%CI: 0.49-1.86)[36,38,42]. No significant difference was observed between the two groups ($P = 0.789$). The between-study heterogeneity was 0 (95%CI: 0-0.9; Supplementary Figure 15). The logHR analysis revealed no significant differences between the groups (Supplementary Figure 16).

Considering the 36-month DFS, the OR was 0.96 (95%CI: 0.4-2.28)[36,38,42]. No significant difference was observed between the two groups ($P = 0.846$). The between-study heterogeneity was 0.05 (95%CI: 0-0.9) was calculated (Supplementary Figure 17). The log HR analysis revealed no significant differences between the groups (Supplemental Figure 18).

Considering the 48-month DFS, the OR was 1.04 (95%CI: 0.31-3.51)[36,38,42]. No significant difference was observed between the two groups ($P = 0.904$). The between-study heterogeneity was 0.32 (95%CI: 0-0.93; Supplementary Figure 19). The log HR analysis revealed no significant differences between the groups (Supplementary Figure 20).

Considering the 60-month DFS, the OR was 1.04 (95%CI: 0.3-3.64)[36,38,42]. No significant difference was observed between the two groups ($P = 0.913$). The between-study heterogeneity was 0.32 (95%CI: 0-0.93) between the groups (Supplementary Figure 21).

Progression-free survival

For the 12-month progression-free survival (PFS), three studies including 340 patients were selected for analysis[40,43,44]. The OR was 0.73 (95%CI: 0.47-1.16). No statistically significant difference was observed between the two groups ($P = 0.101$). The between-study heterogeneity was 0 (95%CI: 0-0.9; Supplementary Figure 22).

Considering the 24-month PFS, the OR was 0.78 (95%CI: 0.1-6.18)[40,43,44]. No significant difference was observed between the two groups ($P = 0.652$). The between-study heterogeneity was 0.72 (95%CI: 0.04-0.92; Supplementary Figure 23).

Considering the 36-month PFS, the OR was 1.04 (95%CI: 0.1-11.05)[40,43,44]. No significant difference was observed between the two groups ($P = 0.946$). The between-study heterogeneity was 0.81 (95%CI: 0.39-0.94; Supplementary Figure 24).

Locoregional recurrence

Data from six studies including 1037 patients were analyzed, revealing locoregional recurrence in 12% of the patients[36,37,41-44]. The OR was 0.98 (95%CI: 0.35-2.77). No significant difference was observed between the two groups ($P = 0.966$). The between-study heterogeneity was 0.76 (95%CI: 0.47-0.89; Supplementary Figure 25).

Distant metastasis recurrence

Data from five studies including 910 patients were analyzed, revealing distal metastasis recurrence in 39% of the patients[37,41-44]. The OR was 1.12 (95%CI: 0.76-1.64). No significant difference was observed between the two groups ($P = 0.462$). The between-study heterogeneity was 0 (95%CI: 0-0.79; Supplementary Figure 26).

Thromboembolism events

Data from four studies including 818 patients were analyzed for the occurrence of thromboembolism events[35,40,42,43]. The OR was 1.93 (95%CI: 0.1-38.65). No significant difference was observed between the two groups ($P = 0.535$). The between-study heterogeneity was 0.72 (95%CI: 0.22-0.90; Supplementary Figure 27).

Cardiac complications

Data from seven studies including 1580 patients were analyzed for the occurrence of cardiac complications[35,36,38,40-43]. The OR was 0.8 (95%CI: 0.42-1.52). No significant difference between the two groups ($P = 0.425$). The between-study heterogeneity was 0.46 (95%CI: 0-0.77; Supplementary Figure 28).

Respiratory complications

Data from seven studies including 1580 patients were analyzed for the occurrence of respiratory complications[35,36,38,40-43]. The OR was 1.04 (95%CI: 0.66-1.64). No significant difference was observed between the two groups ($P = 0.835$). The between-study heterogeneity was 0.59 (95%CI: 0.04-0.82; Supplementary Figure 29).

Renal failure

Data from three studies including 650 patients were analyzed for the occurrence of renal failure[35,42,43]. The OR was 2.43 (95%CI: 1.12-5.28). A statistically significant difference was observed, favoring nCT over nCRT ($P = 0.039$). The between-study heterogeneity was 0 (95%CI: 0-0.9; Supplementary Figure 30).

Neutropenia

Data from three studies including 560 patients were analyzed for the occurrence of neutropenia[35,40,43]. The OR was 0.97 (95%CI: 0.09-10.29). No significant difference was observed between the two groups ($P = 0.964$). The between-study heterogeneity was 0.47 (95%CI: 0-0.84; Supplementary Figure 31).

Anastomotic leakage

Data from seven studies including 1580 patients were analyzed for the occurrence of anastomotic leakage[35,36,38,40-43]. The OR was 0.83 (95%CI: 0.41-1.68). No significant difference was observed between the two groups ($P = 0.539$). The between-study heterogeneity was 0.75 (95%CI: 0.48-0.88; Supplementary Figure 32).

Chyle leakage

Data from six studies including 1366 patients were analyzed for the occurrence of chyle leakage[35,36,40-43]. The OR was 0.99 (95%CI: 0.61-1.61). No significant difference was observed between the two groups ($P = 0.961$). The between-study heterogeneity was 0 (95%CI: 0.48-0.75; Supplementary Figure 33).

Wound infection

Data from five studies including 1022 patients were analyzed for the occurrence of wound infection[35,38,40,42,43]. The OR was 1.04 (95%CI: 0.36-3.02). No significant difference was observed between the two groups ($P = 0.930$). The between-study heterogeneity was 0.37 (95%CI: 0-0.76; Supplementary Figure 34).

Bleeding

Data from four studies including 849 patients were analyzed for the occurrence of bleeding[35,36,40,42]. The OR was 1.4 (95%CI: 0.425-7.79). No statistically significant difference was observed between the two groups ($P = 0.581$). The between-study heterogeneity was 0 (95%CI: 0-0.85; Supplementary Figure 35).

Vocal cord paresis

Data from three studies including 733 patients were analyzed for the occurrence of vocal cord paresis[35,40,42]. The OR was 1.21 (95%CI: 0.04-41.98). No significant difference was

observed between the two groups ($P = 0.537$). The between-study heterogeneity was 0.5 (95%CI: 0-0.85; Supplementary Figure 36).

Leukopenia

Two studies including 485 patients were selected for descriptive analyses[35,40]. Leukopenia occurred in 8% and 12% of the patients in the nCT and nCRT groups, respectively.

Anemia

Two studies including 485 patients were selected for descriptive analyses[35,40]. Anemia occurred in 1% and 0.4% of the patients in the nCT and nCRT groups, respectively.

Nausea or vomiting

Three studies including 560 patients were selected for descriptive analyses[35,40,43]. Nausea or vomiting occurred in 9% and 3% of the patients in the nCT and nCRT groups, respectively.

Diarrhea

Two studies including 485 patients were selected for descriptive analyses[35,40]. Diarrhea occurred in 7% in the nCT group, whereas no cases were noted in the nCRT group.

Hospital stay

Two studies including 430 patients were selected for descriptive analyses[40,42]. The estimated mean hospital stay was 20 (range: 7-97) d in the nCT group and 18 (range: 7-75) d in the nCRT group.

Risk of bias

As we expected, the two RCTs demonstrated a low risk of bias. However, for other included trials, ROB2 indicated some concerns, mainly due to the randomization process (D1). In one trial, concerns were noted regarding the measurement process due to the utilization of a plot digitizer[17]. No instance of high risk of bias was identified across the included studies. Additional information is presented in Table 3.

GRADE approach

Employing the GRADE approach, our findings were determined to have low certainty for most outcomes; moderate certainty for 30-d mortality; very low certainty for 12-month OS, 36-month PFS, and the occurrence of thromboembolism events. The use of RoB2 indicated a moderate risk for all outcomes. High heterogeneity was reported for 36-month PFS and the occurrence of thromboembolic events. Imprecision was observed for pCR and 12-month OS. Additionally, a high variation in oncological treatments decreased the evidence quality, whereas a large effect size increased the quality of pCR and 30-d mortality[49] (Supplementary Table 4).

DISCUSSION

The benefits of neoadjuvant therapy have been previously reported[7-9]. Previous meta-analyses have examined the amplification of nCT and chemoradiotherapy in patients with AC or SCC. In the nCRT group, advantages were observed in terms of 3-year survival with R0 resection; however, the pCR rate had no effect on long-term survival. Perioperative mortality and cardiovascular complications are more common in patients with AC in the nCRT group[50]. A previous network meta-analysis showed that triplet-based chemotherapy increases overall survival and DFS in cases of AC of the stomach or GEJ[51].

pCR is defined as the lack of tumor in the resected specimen or lymph nodes (pT0 pN0 cM0)[15,36]. The 5-year survival rate is presumably 88% in patients with pCR compared to 39% in those without pCR[15,52]. According to a recent investigation comparing the long-term survival of the total population and patients with TRG grade 1-

2 who underwent nCT or chemoradiotherapy before surgery revealed that tumor regression after neoadjuvant treatment is significantly associated with long-term survival, regardless of the treatment regimen[53]. Another retrospective cohort study revealed improved OS and DFS in patients who achieved pCR following nCT compared to those who achieved a lower rate of pCR following nCRT. The authors found a significant association between TRG and survival in both the groups. Additionally, patients who achieved pCR in the nCRT group did not have as good a survival rate as those in the nCT group, although their proportion was higher in the nCRT group. This finding suggests that esophageal AC should be considered a systemic disease and treated accordingly[53,54]. However, other trials have reported that a larger number of patients who achieved pCR do not have improved overall survival[55]. In this meta-analysis, we found a significantly higher pCR in the nCRT group; however, no differences were found in OS, DFS, or PFS, consistent with the findings of previous meta-analyses[50,55]. Based on this finding, we inferred that there is no association between pCR and OS; therefore, the use of pCR as a prognostic factor should be considered in cases of AC. These findings aligned with those of Gebauer *et al's* study[56] reporting that high pCR after CROSS regimen is not clearly associated with longer overall survival[56]. Another study concluded that only clinically complete response without nodal metastasis is associated with long-term survival; therefore, the “watch-and-wait,” strategies should be considered carefully and applied only to patients who have achieved pCR[57]. The utility of pCR as a prognostic indicator of neoadjuvant therapy remains questionable, indicating the need for large number of randomized studies in the future.

Our analysis revealed that none of the investigated groups were superior considering local recurrence, which aligns with the findings of a previous meta-analysis[50]. This indicates that the higher local control provided by radiotherapy does not reduce the incidence of local recurrence. Additionally, we did not detect a significant difference in terms of metastases, which occurred in 39% of the cases compared to 12% of local recurrence cases, suggesting that AC should be treated as a systematic disease, and therefore, the “watch-and-wait” strategies should be considered critically.

Our findings revealed a significantly higher 30-d mortality risk in the chemoradiotherapy group. This can be attributed to complications arising in the postoperative period. However, differences in the outcomes of surgical complications were not noted, consistent with the findings of a previous meta-analysis, in which no difference was reported in anastomotic leakage[50]. Additionally, a previous meta-analysis reported a higher risk of mortality in the postoperative period among patients with AC. Therefore, further investigation into the effects of nCRT on postoperative complications is warranted[50]. We only performed descriptive analysis, which revealed a comparable duration of postoperative hospitalization in both the groups[40,42].

We observed no difference in any of the clinical complications in both the groups; however, a previous meta-analysis reported a higher risk of cardiovascular complications in the nRCT group than in the nCT group, which could be a toxic side effect of this modality.

nCT and radiotherapy are also associated with adverse events, including thromboembolic events, neutropenia, leukopenia, anemia, nausea or vomiting, and diarrhea[35]. Renal failure occurred more often in the nCT group than in the nCRT group, indicating a toxic side effect of nCT. However, no difference was reported in terms of cardiac failure, in contrast to a former meta-analysis[50]. According to previous investigations, neutropenia is not associated with either neoadjuvant treatment modality. In the descriptive analysis, leukopenia occurred 4% more frequently in the nCRT group than in the nCT group, making them more vulnerable to developing infections. Additionally, a low number of anemia cases was observed in both the groups. The quality of life can be assessed using the EORTC QLQ-C30 questionnaire[58], which includes encompasses side effects including nausea, vomiting, and diarrhea. Notably, these side effects occurred approximately 7% more frequently in the nCT group than in the nCRT group.

Our meta-analysis provides the most comprehensive and recent summary of the data, particularly focusing on patients with esophageal AC. In addition, various outcomes were analyzed in a sufficient number of patients. The data from this study

accurately reflect the esophageal AC population. No significant differences in demographic characteristics were reported between patients of the two groups.

Nevertheless, our study has some limitations. Deviating from the protocol, we included propensity score-matched studies and cohort trials, which are less reliable than RCTs and have potentially significant biases. Additionally, all trials were conducted in Western countries, reflecting a characteristic of AC, thus limiting the generalizability of the results to the Asian population or other countries. The use of various neoadjuvant regimens and some the lack of separation between preoperative and perioperative therapies in some included studies also pose some limitations. Furthermore, the evidence for most outcomes was deemed low; therefore, the true effect may differ substantially from the estimate.

In summary, one might question the lack of impact of radiotherapy on overall survival, despite improvements in measures of pathological regression, known to correlate with survival. This discrepancy can be attributed to modification of these crucial measures by local therapy. In the context of modern surgical techniques, the systemic component of the disease is the primary determinant of survival in esophageal and gastroesophageal junction ACs. Hence, the incorporation of systemic chemotherapy, new immunotherapies, and targeted treatments capable of addressing distant diseases holds greater potential to enhance patient survival in the future.

CONCLUSION

In patients with esophageal AC, neoadjuvant chemoradiation increases pCR and 30-d mortality; however, it has no effect on long-term survival. nCT may be associated with side effects that can decrease the quality of life. Further randomized trials are required to address the limitations in the quality of the available studies.

ACKNOWLEDGEMENTS

REFERENCES

- 1 **Liu CQ**, Ma YL, Qin Q, Wang PH, Luo Y, Xu PF, Cui Y. Epidemiology of esophageal cancer in 2020 and projections to 2030 and 2040. *Thorac Cancer* 2023; **14**: 3-11 [PMID: 36482832 DOI: 10.1111/1759-7714.14745]
- 2 **Tinusz B**, Szapáry LB, Paládi B, Papp A, Bogner B, Hegedűs I, Bellyei S, Vincze Á, Solt J, Micsik T, Dunás-Varga V, Pályu E, Vass T, Schnabel T, Farkas N, Hegyi P, Thrift AP, Erőss B. The Esophageal Adenocarcinoma Epidemic Has Reached Hungary: A Multicenter, Cross-Sectional Study. *Front Oncol* 2020; **10**: 541794 [PMID: 33425714 DOI: 10.3389/fonc.2020.541794]
- 3 **Rice TW**, Patil DT, Blackstone EH. 8th edition AJCC/UICC staging of cancers of the esophagus and esophagogastric junction: application to clinical practice. *Ann Cardiothorac Surg* 2017; **6**: 119-130 [PMID: 28447000 DOI: 10.21037/acs.2017.03.14]
- 4 **Liu F**, Wang W, Wang C, Peng X. Enhanced recovery after surgery (ERAS) programs for esophagectomy protocol for a systematic review and meta-analysis. *Medicine (Baltimore)* 2018; **97**: e0016 [PMID: 29465538 DOI: 10.1097/MD.00000000000010016]
- 5 **Sindler DL**, Mátrai P, Szakó L, Berki D, Berke G, Csontos A, Papp C, Hegyi P, Papp A. Faster recovery and bowel movement after early oral feeding compared to late oral feeding after upper GI tumor resections: a meta-analysis. *Front Surg* 2023; **10**: 1092303 [PMID: 37304183 DOI: 10.3389/fsurg.2023.1092303]
- 6 **Szakó L**, Németh D, Farkas N, Kiss S, Dömötör RZ, Engh MA, Hegyi P, Eross B, Papp A. Network meta-analysis of randomized controlled trials on esophagectomies in esophageal cancer: The superiority of minimally invasive surgery. *World J Gastroenterol* 2022; **28**: 4201-4210 [PMID: 36157121 DOI: 10.3748/wjg.v28.i30.4201]
- 7 **van Hagen P**, Hulshof MC, van Lanschot JJ, Steyerberg EW, van Berge Henegouwen MI, Wijnhoven BP, Richel DJ, Nieuwenhuijzen GA, Hospers GA, Bonenkamp JJ, Cuesta MA, Blaisse RJ, Busch OR, ten Kate FJ, Creemers GJ, Punt CJ, Plukker JT, Verheul HM, Spillenaar Bilgen EJ, van Dekken H, van der Sangen MJ, Rozema T, Biermann K, Beukema JC, Piet AH, van Rij CM, Reinders JG, Tilanus HW, van der Gaast A; CROSS Group.

Preoperative chemoradiotherapy for esophageal or junctional cancer. *N Engl J Med* 2012; **366**: 2074-2084 [PMID: 22646630 DOI: 10.1056/NEJMoa1112088]

8 **Cunningham D**, Allum WH, Stenning SP, Thompson JN, Van de Velde CJ, Nicolson M, Scarffe JH, Lofts FJ, Falk SJ, Iveson TJ, Smith DB, Langley RE, Verma M, Weeden S, Chua YJ, MAGIC Trial Participants. Perioperative chemotherapy *vs* surgery alone for resectable gastroesophageal cancer. *N Engl J Med* 2006; **355**: 11-20 [PMID: 16822992 DOI: 10.1056/NEJMoa055531]

9 **Ychou M**, Boige V, Pignon JP, Conroy T, Bouché O, Lebreton G, Ducourtieux M, Bedenne L, Fabre JM, Saint-Aubert B, Genève J, Lasser P, Rougier P. Perioperative chemotherapy compared with surgery alone for resectable gastroesophageal adenocarcinoma: an FNCLCC and FFCD multicenter phase III trial. *J Clin Oncol* 2011; **29**: 1715-1721 [PMID: 21444866 DOI: 10.1200/JCO.2010.33.0597]

10 **Allum WH**, Stenning SP, Bancewicz J, Clark PI, Langley RE. Long-term results of a randomized trial of surgery with or without preoperative chemotherapy in esophageal cancer. *J Clin Oncol* 2009; **27**: 5062-5067 [PMID: 19770374 DOI: 10.1200/JCO.2009.22.2083]

11 **Ronellenfitsch U**, Schwarzbach M, Hofheinz R, Kienle P, Kieser M, Slinger TE, Burmeister B, Kelsen D, Niedzwiecki D, Schuhmacher C, Urba S, van de Velde C, Walsh TN, Ychou M, Jensen K. Preoperative chemo(radio)therapy *vs* primary surgery for gastroesophageal adenocarcinoma: systematic review with meta-analysis combining individual patient and aggregate data. *Eur J Cancer* 2013; **49**: 3149-3158 [PMID: 23800671 DOI: 10.1016/j.ejca.2013.05.029]

12 **Ronellenfitsch U**, Schwarzbach M, Hofheinz R, Kienle P, Kieser M, Slinger TE, Jensen K; GE Adenocarcinoma Meta - analysis Group. Perioperative chemo(radio)therapy *vs* primary surgery for resectable adenocarcinoma of the stomach, gastroesophageal junction, and lower esophagus. *Cochrane Database Syst Rev* 2013: CD008107 [PMID: 23728671 DOI: 10.1002/14651858.CD008107.pub2]

13 **Schuhmacher C**, Gretscher S, Lordick F, Reichardt P, Hohenberger W, Eisenberger CF, Haag C, Mauer ME, Hasan B, Welch J, Ott K, Hoelscher A, Schneider PM, Bechstein W, Wilke H, Lutz MP, Nordlinger B, Van Cutsem E, Siewert JR, Schlag PM. Neoadjuvant

chemotherapy compared with surgery alone for locally advanced cancer of the stomach and cardia: European Organisation for Research and Treatment of Cancer randomized trial 40954. *J Clin Oncol* 2010; **28**: 5210-5218 [PMID: 21060024 DOI: 10.1200/JCO.2009.26.6114]

14 **Sjoquist KM**, Burmeister BH, Smithers BM, Zalcberg JR, Simes RJ, Barbour A, Gebiski V; Australasian Gastro-Intestinal Trials Group. Survival after neoadjuvant chemotherapy or chemoradiotherapy for resectable oesophageal carcinoma: an updated meta-analysis. *Lancet Oncol* 2011; **12**: 681-692 [PMID: 21684205 DOI: 10.1016/S1470-2045(11)70142-5]

15 **Stahl M**, Walz MK, Stuschke M, Lehmann N, Meyer HJ, Riera-Knorrenschild J, Langer P, Engenhart-Cabillic R, Bitzer M, Königsrainer A, Budach W, Wilke H. Phase III comparison of preoperative chemotherapy compared with chemoradiotherapy in patients with locally advanced adenocarcinoma of the esophagogastric junction. *J Clin Oncol* 2009; **27**: 851-856 [PMID: 19139439 DOI: 10.1200/JCO.2008.17.0506]

16 **National Institute for Health and Care Research**. PROSPERO is fast-tracking registration of protocols related to COVID-19. [cited 23 February 2024]. Available from: <https://www.crd.york.ac.uk/prospero/>

17 **Plotdigitizer**. All-in-One Tool to Extract Data from Graphs, Plots & Images. [cited 23 February 2024]. Available from: <https://plotdigitizer.com/app>

18 **Guyot P**, Ades AE, Ouwers MJ, Welton NJ. Enhanced secondary analysis of survival data: reconstructing the data from published Kaplan-Meier survival curves. *BMC Med Res Methodol* 2012; **12**: 9 [PMID: 22297116 DOI: 10.1186/1471-2288-12-9]

19 **WebPlotDigitizer**. Extract data from XY charts, Bar graphs, Polar diagrams and much more! [cited 23 February 2024]. Available from: <https://automeris.io/WebPlotDigitizer/>

20 **Higgins JP**, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; **21**: 1539-1558 [PMID: 12111919 DOI: 10.1002/sim.1186]

21 **Harbord RM**, Harris RJ, Sterne JAC. Updated tests for small-study effects in meta-analyses. *The Stata Journal: Promoting Communications on Statistics and Stata* 2009; **9**: 197-210 [DOI: 10.1177/1536867X0900900202]

- 22 **Harrer M**, Cuijpers P, Toshi F, Ebert DD. Doing meta-analysis with R: a hands-on guide. 1st ed. New York: Chapman and Hall/CRC Press, 2021 [DOI: 10.1201/9781003107347]
- 23 **R Core Team**. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. 2023. [cited 23 February 2024]. Available from: <https://www.R-project.org/>
- 24 **Schwarzer G**. Meta-Analysis in R. In: Systematic Reviews in Health Research: Meta - Analysis in Context, 3rd Ed. Egger M, Higgins JPT, Smith GD, editors. New York: John Wiley & Sons, 2022 [DOI: 10.1002/9781119099369.ch26]
- 25 **Cuijpers P**, Furukawa T, Ebert DD. Dmetar: companion R package for the guide doing meta-analysis in R. 2022. [cited 23 February 2024]. Available from: <https://dmetar.protectlab.org>
- 26 **Mantel N**, Haenszel W. Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst* 1959; **22**: 719-748 [PMID: 13655060]
- 27 **Robins J**, Greenland S, Breslow NE. A general estimator for the variance of the Mantel-Haenszel odds ratio. *Am J Epidemiol* 1986; **124**: 719-723 [PMID: 3766505 DOI: 10.1093/oxfordjournals.aje.a114447]
- 28 **Cooper HM**, Hedges LV, Valentine JC, editors. The handbook of research synthesis and meta-analysis. 2nd ed. New York: Russell Sage Foundation, 2009
- 29 **Sweeting MJ**, Sutton AJ, Lambert PC. What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data. *Stat Med* 2004; **23**: 1351-1375 [PMID: 15116347 DOI: 10.1002/sim.1761]
- 30 **Knapp G**, Hartung J. Improved tests for a random effects meta-regression with a single covariate. *Stat Med* 2003; **22**: 2693-2710 [PMID: 12939780 DOI: 10.1002/sim.1482]
- 31 **IntHout J**, Ioannidis JP, Borm GF. The Hartung-Knapp-Sidik-Jonkman method for random effects meta-analysis is straightforward and considerably outperforms the standard DerSimonian-Laird method. *BMC Med Res Methodol* 2014; **14**: 25 [PMID: 24548571 DOI: 10.1186/1471-2288-14-25]

- 32 **Paule RC**, Mandel J. Consensus Values and Weighting Factors. *J Res Natl Bur Stand* (1977) 1982; **87**: 377-385 [PMID: 34566088 DOI: 10.6028/jres.087.022]
- 33 **Veroniki AA**, Jackson D, Viechtbauer W, Bender R, Bowden J, Knapp G, Kuss O, Higgins JP, Langan D, Salanti G. Methods to estimate the between-study variance and its uncertainty in meta-analysis. *Res Synth Methods* 2016; **7**: 55-79 [PMID: 26332144 DOI: 10.1002/jrsm.1164]
- 34 **Shinyapps**. Estimating the sample mean and standard deviation. [cited 23 February 2024]. Available from: <https://smcgrath.shinyapps.io/estmeansd>
- 35 **Anderegg MCJ**, van der Sluis PC, Ruurda JP, Gisbertz SS, Hulshof MCCM, van Vulpen M, Mohammed NH, van Laarhoven HWM, Wiezer MJ, Los M, van Berge Henegouwen MI, van Hillegersberg R. Preoperative Chemoradiotherapy Versus Perioperative Chemotherapy for Patients With Resectable Esophageal or Gastroesophageal Junction Adenocarcinoma. *Ann Surg Oncol* 2017; **24**: 2282-2290 [PMID: 28424936 DOI: 10.1245/s10434-017-5827-1]
- 36 **Luc G**, Vendrely V, Terrebonne E, Chiche L, Collet D. Neoadjuvant chemoradiotherapy improves histological results compared with perioperative chemotherapy in locally advanced esophageal adenocarcinoma. *Ann Surg Oncol* 2015; **22**: 604-609 [PMID: 25169119 DOI: 10.1245/s10434-014-4005-y]
- 37 **Münch S**, Habermehl D, Agha A, Belka C, Combs SE, Eckel R, Friess H, Gerbes A, Nüssler NC, Schepp W, Schmid RM, Schmitt W, Schubert-Fritschle G, Weber B, Werner J, Engel J. Perioperative chemotherapy vs. neoadjuvant chemoradiation in gastroesophageal junction adenocarcinoma: A population-based evaluation of the Munich Cancer Registry. *Strahlenther Onkol* 2018; **194**: 125-135 [PMID: 29071366 DOI: 10.1007/s00066-017-1225-7]
- 38 **Spicer JD**, Stiles BM, Sudarshan M, Correa AM, Ferri LE, Altorki NK, Hofstetter WL. Preoperative Chemoradiation Therapy Versus Chemotherapy in Patients Undergoing Modified En Bloc Esophagectomy for Locally Advanced Esophageal Adenocarcinoma: Is Radiotherapy Beneficial? *Ann Thorac Surg* 2016; **101**: 1262-9; discussion 1969-70 [PMID: 26916717 DOI: 10.1016/j.athoracsur.2015.11.070]

- 39 **Favi F**, Bollschweiler E, Berlth F, Plum P, Hescheler DA, Alakus H, Semrau R, Celik E, Mönig SP, Drebber U, Hölscher AH. Neoadjuvant chemotherapy or chemoradiation for patients with advanced adenocarcinoma of the oesophagus? A propensity score-matched study. *Eur J Surg Oncol* 2017; **43**: 1572-1580 [PMID: 28666624 DOI: 10.1016/j.ejso.2017.06.003]
- 40 **Goense L**, van der Sluis PC, van Rossum PSN, van der Horst S, Meijer GJ, Haj Mohammad N, van Vulpen M, Mook S, Ruurda JP, van Hillegersberg R. Perioperative chemotherapy *vs* neoadjuvant chemoradiotherapy for esophageal or GEJ adenocarcinoma: A propensity score-matched analysis comparing toxicity, pathologic outcome, and survival. *J Surg Oncol* 2017; **115**: 812-820 [PMID: 28267212 DOI: 10.1002/jso.24596]
- 41 **Markar SR**, Noordman BJ, Mackenzie H, Findlay JM, Boshier PR, Ni M, Steyerberg EW, van der Gaast A, Hulshof MCCM, Maynard N, van Berge Henegouwen MI, Wijnhoven BPL, Reynolds JV, Van Lanschot JJB, Hanna GB. Multimodality treatment for esophageal adenocarcinoma: multi-center propensity-score matched study. *Ann Oncol* 2017; **28**: 519-527 [PMID: 28039180 DOI: 10.1093/annonc/mdw560]
- 42 **Visser E**, Edholm D, Smithers BM, Thomson IG, Burmeister BH, Walpole ET, Gotley DC, Joubert WL, Atkinson V, Mai T, Thomas JM, Barbour AP. Neoadjuvant chemotherapy or chemoradiotherapy for adenocarcinoma of the esophagus. *J Surg Oncol* 2018; **117**: 1687-1696 [PMID: 29806960 DOI: 10.1002/jso.25089]
- 43 **Burmeister BH**, Thomas JM, Burmeister EA, Walpole ET, Harvey JA, Thomson DB, Barbour AP, Gotley DC, Smithers BM. Is concurrent radiation therapy required in patients receiving preoperative chemotherapy for adenocarcinoma of the oesophagus? A randomised phase II trial. *Eur J Cancer* 2011; **47**: 354-360 [PMID: 21084184 DOI: 10.1016/j.ejca.2010.09.009]
- 44 **Stahl M**, Walz MK, Riera-Knorrenschild J, Stuschke M, Sandermann A, Bitzer M, Wilke H, Budach W. Preoperative chemotherapy *vs* chemoradiotherapy in locally advanced adenocarcinomas of the oesophagogastric junction (POET): Long-term results

of a controlled randomised trial. *Eur J Cancer* 2017; **81**: 183-190 [PMID: 28628843 DOI: 10.1016/j.ejca.2017.04.027]

45 **Alderson D**, Langley RE, Nankivell MG, Blazeby JM, Griffin M, Crellin A, Grabsch HI, Okines AFC, Goldstein C, Falk S, Thompson J, Krysztopik R, Coxon FY, Pritchard S, Langer R, Stenning SP, Cunningham D. Neoadjuvant chemotherapy for resectable oesophageal and junctional adenocarcinoma: results from the UK Medical Research Council randomised OEO5 trial (ISRCTN 01852072). *J Clin Oncol* 2015; **33**: Abstr 4002 [DOI: 10.1200/jco.2015.33.15_suppl.4002]

46 **Shapiro J**, van Lanschot JJB, Hulshof MCCM, van Hagen P, van Berge Henegouwen MI, Wijnhoven BPL, van Laarhoven HWM, Nieuwenhuijzen GAP, Hospers GAP, Bonenkamp JJ, Cuesta MA, Blaisse RJB, Busch ORC, Ten Kate FJW, Creemers GM, Punt CJA, Plukker JTM, Verheul HMW, Bilgen EJS, van Dekken H, van der Sangen MJC, Rozema T, Biermann K, Beukema JC, Piet AHM, van Rij CM, Reinders JG, Tilanus HW, Steyerberg EW, van der Gaast A; CROSS study group. Neoadjuvant chemoradiotherapy plus surgery *vs* surgery alone for oesophageal or junctional cancer (CROSS): long-term results of a randomised controlled trial. *Lancet Oncol* 2015; **16**: 1090-1098 [PMID: 26254683 DOI: 10.1016/S1470-2045(15)00040-6]

47 **Lorenzen S**, Pauligk C, Homann N, Schmalenberg H, Jäger E, Al-Batran SE. Feasibility of perioperative chemotherapy with infusional 5-FU, leucovorin, and oxaliplatin with (FLOT) or without (FLO) docetaxel in elderly patients with locally advanced esophagogastric cancer. *Br J Cancer* 2013; **108**: 519-526 [PMID: 23322206 DOI: 10.1038/bjc.2012.588]

48 **Ferri LE**, Ades S, Alcindor T, Chasen M, Marcus V, Hickeson M, Artho G, Thirlwell MP. Perioperative docetaxel, cisplatin, and 5-fluorouracil (DCF) for locally advanced esophageal and gastric adenocarcinoma: a multicenter phase II trial. *Ann Oncol* 2012; **23**: 1512-1517 [PMID: 22039085 DOI: 10.1093/annonc/mdr465]

49 **Kirmayr M**, Quilodrán C, Valente B, Loezar C, Garegnani L, Franco JVA. The GRADE approach, Part 1: how to assess the certainty of the evidence. *Medwave* 2021; **21**: e8109 [PMID: 33830974 DOI: 10.5867/medwave.2021.02.8109]

50 **Han J**, Wang Z, Liu C. Survival and complications after neoadjuvant chemotherapy or chemoradiotherapy for esophageal cancer: a meta-analysis. *Future Oncol* 2021; **17**: 2257-2274 [PMID: 33739165 DOI: 10.2217/fon-2021-0021]

51 **Grizzi G**, Petrelli F, Di Bartolomeo M, Viti M, Texeira Moraes M, Luciani A, Passalacqua R, Ghidini M, Tomasello G, Baiocchi GL, Celotti A. Preferred neoadjuvant therapy for gastric and gastroesophageal junction adenocarcinoma: a systematic review and network meta-analysis. *Gastric Cancer* 2022; **25**: 982-987 [PMID: 35704113 DOI: 10.1007/s10120-022-01314-9]

52 **Barbour AP**, Jones M, Gonen M, Gotley DC, Thomas J, Thomson DB, Burmeister B, Smithers BM. Refining esophageal cancer staging after neoadjuvant therapy: importance of treatment response. *Ann Surg Oncol* 2008; **15**: 2894-2902 [PMID: 18663531 DOI: 10.1245/s10434-008-0084-y]

53 **Sciuto M**, Capovilla G, Scarton A, Tagkalos E, Uzun E, Moletta L, Hadzijusufović Edin, Provenzano L, Salvador R, Pierobon E, Zanchettin G, Berlth F, Lang H, Grimminger P, Valmasoni M. 462. Major pathologic response in esophageal adenocarcinoma: should we adopt a new paradigm in defining response to treatments?. *Diseases of the Esophagus* 2003; **36** [DOI: 10.1093/dote/doad052.246]

54 **Scarton A**, Capovilla G, Tagkalos E, Uzun E, Moletta L, Hadzijusufović E, Provenzano L, Salvador R, Pierobon E, Zanchettin G, Berlth F, Grimminger P, Valmasoni M. 463. The impact of pathological tumor response following neoadjuvant chemotherapy and chemoradiotherapy for esophageal adenocarcinoma. A retrospective multicenter cohort study. *Diseases of the Esophagus* 2003; **36** [DOI: 10.1093/dote/doad052.247]

55 **Klevebro F**, Alexandersson von Döbeln G, Wang N, Johnsen G, Jacobsen AB, Friesland S, Hatlevoll I, Glenjen NI, Lind P, Tsai JA, Lundell L, Nilsson M. A randomized clinical trial of neoadjuvant chemotherapy *vs* neoadjuvant chemoradiotherapy for cancer of the oesophagus or gastro-oesophageal junction. *Ann Oncol* 2016; **27**: 660-667 [PMID: 26782957 DOI: 10.1093/annonc/mdw010]

56 **Gebauer F**, Plum PS, Damanakis A, Chon SH, Popp F, Zander T, Quaas A, Fuchs H, Schmidt T, Schröder W, Bruns CJ. Long-Term Postsurgical Outcomes of Neoadjuvant

Chemoradiation (CROSS) Versus Chemotherapy (FLOT) for Multimodal Treatment of Adenocarcinoma of the Esophagus and the Esophagogastric Junction. *Ann Surg Oncol* 2023; **30**: 7422-7433 [PMID: 37210683 DOI: 10.1245/s10434-023-13643-9]

57 **Schroeder W**, Ghadimi MPH, Schloesser H, Loeser H, Schiller P, Zander T, Gebauer F, Fuchs H, Quaas A, Bruns CJ. Long-Term Outcome After Histopathological Complete Response with and Without Nodal Metastases Following Multimodal Treatment of Esophageal Cancer. *Ann Surg Oncol* 2022 [PMID: 35403919 DOI: 10.1245/s10434-022-11700-3]

58 **Adenis A**, Kulkarni AS, Girotto GC, de la Fouchardiere C, Senellart H, van Laarhoven HWM, Mansoor W, Al-Rajabi R, Norquist J, Amonkar M, Suryawanshi S, Bhagia P, Metges JP. Impact of Pembrolizumab Versus Chemotherapy as Second-Line Therapy for Advanced Esophageal Cancer on Health-Related Quality of Life in KEYNOTE-181. *J Clin Oncol* 2022; **40**: 382-391 [PMID: 34730989 DOI: 10.1200/JCO.21.00601]

Footnotes

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Corresponding Author's Membership in Professional Societies:

Specialty type: Gastroenterology and hepatology

Country/Territory of origin: Hungary

Peer-review report's classification

Scientific Quality: Grade A

Novelty:

Creativity or Innovation:

Scientific Significance:

P-Reviewer: S-Editor: L-Editor: P-Editor:

Figures and Tables




Figure 1 The preferred reporting items for systematic reviews and meta-analyses flow diagram flowchart shows the number of articles (n) in the different selection stages of the selection process. AC: Adenocarcinoma; SCC: Squamous cell carcinoma.




Figure 2 Analysis of pathological complete response. OR: Odds ratio.




Figure 3 Analysis of the 30-d mortality. nCT: Neoadjuvant chemotherapy; nCRT: Neoadjuvant chemoradiotherapy; OR: Odds ratio.




Figure 4 The Kaplan-Meier curves for the overall survival. The x-axis shows the time in month, the y-axis shows the number of patients in percentage. nCT: Neoadjuvant chemotherapy; nCRT: Neoadjuvant chemoradiotherapy; PFS: Progression-free survival.




Figure 5 Pooled hazard ratio analysis of the overall mortality. HR: Hazard ratio; HK: Hoffman-Kringle random effect model.

Table 1 Characteristic of the studies

Ref.	Design	Center	Country	Year	Number of patients	nCT	nCRT	AC, %
	RCT, Phase							
Stahl <i>et al</i> [44], 2017	III	1	Germany	N/A	119	59	60	100
Burmeister <i>et al</i> [43], 2011	RCT, Phase II	1	Australia	N/A	75	36	39	100
Visser <i>et al</i> [42], 2018	PSM	1	Australia	2000-2017	262	131	131	100
Markar <i>et al</i> [41], 2017	PSM	10	United Kingdom	2001-2012	442	221	221	100
Goense <i>et al</i> [40], 2017	PSM	1	Netherlands	2006-2015	172	86	86	100
Favi <i>et al</i> [39], 2017	PSM	1	Germany	2011-2015	80	40	40	100
Anderegg <i>et al</i> [35], 2017	Cohort, PID 3	3	Netherlands	2005-2011	313	137	176	100
Spicer <i>et al</i> [38], 2016	Cohort, PID 3	3	United States	2002-2012	214	114	100	100
Luc <i>et al</i> [36], 2015	Cohort, PID 1	1	France	2000-2012	116	61	55	100
	Cohort,							
Münch <i>et al</i> [37], 2018	PCCR	70	Germany	1998-2014	135	71	64	100

RCT: Randomized controlled trial; PSM: Propensity score matched cohort; PID: Prospective institutional databases; PCCR: Population-based clinical cancer registry; N/A: Not applicable; nCT: Neoadjuvant chemotherapy; nCRT: Neoadjuvant chemoradiotherapy; AC: Adenocarcinoma.

Table 2 Characteristic of the neoadjuvant regimen, *n* (%)

Ref.	nCT	nCRT	
	Chemotherapy	Chemotherapy	Irradiation
Stahl <i>et al</i> [44], 2017	15 × weekly CFFa	15 × weekly CFFa	30 Gy in 15
		followed by 3 wk fractions of 2 Gy	
Burmeister <i>et al</i> [43], 2011		course of CRT + 1 in 3 wk	
		cycle CE	
	C (80 mg/m ²) + iv 5-FU	CF + RT, 5-FU	35 Gy in 15
	(1000 mg/m ² /d) on	reduced to 800	fractions in 3 wk
	days 1 and 21	mg/m ² /d (on day 21)	(on day 21)
	OEO2	OEO2 + RT	35 Gy in 15
			fractions or 45
			Gy in 25
			fractions
	MAGIC	DCF (2 cycles pre-	45 Gy in 25
		operatively) + RT	fractions
	DCF (2 cycles pre-	CROSS (since 2015)	41.4 Gy in 23
	operatively)		fractions of 1.8
			Gy in 5 wk
	Cisplatin + 5-FU: 92 (70)	Cisplatin + 5-FU: 94	35 Gy: 69 (53)
		(72)	
Visser <i>et al</i> [42], 2018	Epirubicin, cisplatin, 5-	Epirubicin, cisplatin,	41 Gy: 14 (11)
	FU: 30 (23)	5-FU: 2 (2)	
	Carboplatin + paclitaxel:	Carboplatin +	45 Gy: 40 (31)
	0 (0)	paclitaxel: 20 (15)	
	Other: 9 (7)	Other: 15 (11)	Other: 8 (6)

Markar <i>et al</i> [41], 2017	mainly MAGIC, OEO2 or OEO5 regimens[8,10,45]	CROSS regimen[7,46]	41.4 Gy in 23 fractions of 1.8 Gy in 5 wk
Goense <i>et al</i> [40], 2017	ECX	CROSS regimen[7,46]	41.4 Gy in 23 fractions of 1.8 Gy in 5 wk
Favi <i>et al</i> [39] , 2017	FLOT[47]	CROSS regimen[7,46]	41.4 Gy in 23 fractions of 1.8 Gy in 5 wk
Anderegg <i>et al</i> [35], 2017	ECX	CROSS regimen[7,46]	41.4 Gy in 23 fractions of 1.8 Gy in 5 wk
Spicer <i>et al</i> [38], 2016	Cornell: Platinum or taxane-based doublet, or both McGill: DCF (3 cycles)[48]	concurrent ChT + RT	50.4 Gy
Luc <i>et al</i> [36], 2015	DCF (3 cycles pre- and postoperatively)	continuous iv 5-FU 750 mg/m ² /d on week days 1–5 by, C 20 mg/m ² on day 1	45 Gy for 5 d per week at 1.8 Gy/d (started on day 28 along with the second CT cycle)
Münch <i>et al</i> [37], 2018	N/A	N/A	N/A

OEO2: Two cycles of cisplatin and 5-fluoruracil (5-FU). OEO5: Five cycles of cisplatin and 5-FU. CFFa: 15 × weekly 5-FU (2000 mg/m², 24 h infusion)/Fa (500 mg/m², 2 h infusion) and biweekly cisplatin (50 mg/m², 1 h infusion), in 14 wk. MAGIC: Protocol epirubicin, cisplatin, and 5-FU for three cycles before and three cycles after esophagectomy (EGJ

Siewert II EAC and good vital status). DCF: Docetaxel 75 mg/m² (on day 1), cisplatin 75 mg/m² (on day 1), and 5-FU 750 mg/m²/d by continuous infusion on days 2–5, (day 1 = day 22 = day 43). CROSS: With concurrent weekly administration of carboplatin (targeted at an area under the curve of 2 mg/mL per min) and paclitaxel (50 mg/m² of body-surface area). Forty-one four Gy in 23 fractions of 1.8 Gy in 5 wk. FLOT: 5-FU/leucovorin, oxaliplatin, and docetaxel 50 mg/m² every 2 wk. ECX: Pre- and postoperative 3-wk cycles epirubicin (50 mg/m²) and cisplatin (60 mg/m²), followed by 1000 mg/m² of capecitabine twice daily for 14 d or 625 mg/m² of capecitabine twice daily for 21 d. Adaptations to the regimen such as dose reduction or change of regimen to oxaliplatin or 5-FU were applied when necessary. C: Cisplatin; F: Fluorouracil; P: Paclitaxel; Fa: Folinic acid; E: Etoposide; N/A: Not applicable; nCT: Neoadjuvant chemotherapy; nCRT: Neoadjuvant chemoradiotherapy; AC: Adenocarcinoma; CRT: Chemoradiotherapy; CT: Chemotherapy; RT: Radiotherapy; 5-FU: 5-fluoruracil.

Table 3 Results using the risk of bias tool 2

Ref.	D1	D2	D3	D4	D5	Overall
Stahl <i>et al</i> [44], 2017	+	+	+	+	+	+
Burmeister <i>et al</i> [43], 2011	+	+	+	+	+	+
Visser <i>et al</i> [42], 2018	!	+	+	+	+	!
Markar <i>et al</i> [41], 2017	!	+	+	+	+	!
Goense <i>et al</i> [40], 2017	!	+	+	+	+	!
Favi <i>et al</i> [39], 2017	!	+	+	+	+	!
Anderegg <i>et al</i> [35], 2017	!	+	+	+	+	!
Spicer <i>et al</i> [38], 2016	!	+	+	+	+	!
Luc <i>et al</i> [36], 2015	!	+	+	+	+	!

Münch *et al*[37], 2018 ! + + ! + !

+: Low risk; !: Some concerns; -: High risk. D1: Randomization process; D2: Deviation from the intended interventions; D3: Missing outcome data; D4: Measurement of the outcome; D5: Selection of the reported result.

Source: Csontos A, Fazekas A, Szakó L, Farkas N, Papp C, Ferenczi S, Bellyei S, Hegyi P, Papp A. Effects of neoadjuvant chemotherapy vs chemoradiotherapy in the treatment of esophageal adenocarcinoma: A systematic review and meta-analysis. *World J Gastroenterol* 2024; 30(11): 1621-1635

URL: <https://www.wjgnet.com/1007-9327/full/v30/i11/1621.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v30.i11.1621>